arXiv:1609.03006v2 [math.AT] 6 Mar 2017

THE SQUARING OPERARTION AND THE SINGER ALGEBRAIC TRANSFER

NGUYỄN SUM

Abstract. Let P_k be the graded polynomial algebra $\mathbb{F}_2[x_1, x_2, \ldots, x_k]$, with the degree of each x_i being 1, regarded as a module over the mod-2 Steenrod algebra $\mathcal{A}$, and let GL_k be the general linear group over the prime field $\mathbb{F}_2$ which acts regularly on P_k. We study the algebraic transfer constructed by Singer using the technique of the *hit problem*. This transfer is a homomorphism from the homology of the mod-2 Steenrod algebra, $\mathrm{Tor}^{\mathcal{A}}_{k,k+d}(\mathbb{F}_2, \mathbb{F}_2)$, to the subspace of $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k$ consisting of all the GL_k-invariant classes of degree d.

In this paper, we extend a result of Hưng in [10] on the relation between the Singer algebraic transfer and the classical squaring operation on the cohomology of the Steenrod algebra. Using this result, we show that Singer's conjecture for the algebraic transfer is true in the case $k = 5$ and the degree $5(2^s - 1)$ with s an arbitrary positive integer.

1. Introduction

Let V_k be an elementary abelian 2-group of rank k and let BV_k be the classifying space of V_k. Then,

$$P_k := H^*(BV_k) \cong \mathbb{F}_2[x_1, x_2, \ldots, x_k],$$

a polynomial algebra in k generators $x_1, x_2, \ldots, x_k$, each of degree 1. Here the cohomology is taken with coefficients in the prime field $\mathbb{F}_2$ of two elements.

Being the cohomology of a topological space, P_k is a module over the mod-2 Steenrod algebra, $\mathcal{A}$. The action of $\mathcal{A}$ on P_k is determined by the elementary properties of the Steenrod squares Sq^i and subject to the Cartan formula (see Steenrod and Epstein [21]).

Let GL_k be the general linear group over the field $\mathbb{F}_2$. Since V_k is an $\mathbb{F}_2$-vector space of dimension k, this group acts naturally on V_k and therefore on the cohomology of BV_k. The two actions of $\mathcal{A}$ and GL_k upon P_k commute with each other. Hence, there is an inherited action of GL_k on $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k$.

For a non-negative integer d, denote by $(P_k)_d$ the subspace of P_k consisting of all the homogeneous polynomials of degree d in P_k and by $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_d$ the subspace of $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k$ consisting of all the classes represented by the elements in $(P_k)_d$. In [19], Singer defined the algebraic transfer, which is a homomorphism

$$\varphi_k : \mathrm{Tor}^{\mathcal{A}}_{k,k+d}(\mathbb{F}_2, \mathbb{F}_2) \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)^{GL_k}_d$$

[1]2010 *Mathematics Subject Classification.* Primary 55S10; 55S05, 55T15.

[2]*Keywords and phrases:* Steenrod algebra, Peterson hit problem, algebraic transfer, polynomial algebra.

[3]The work is funded by the National Foundation for Science and Technology Development (NAFOSTED) of Vietnam under the grant number 101.04-2017.05.

from the homology of the Steenrod algebra, $\mathrm{Tor}^{\mathcal{A}}_{k,k+d}(\mathbb{F}_2,\mathbb{F}_2)$, to the subspace of $(\mathbb{F}_2\otimes_{\mathcal{A}}P_k)_d$ consisting of all the GL_k-invariant classes. The Singer algebraic transfer is a useful tool in describing the homology groups of the Steenrod algebra. It was studied by many authors (see Boardman [1], Bruner-Hà-Hưng [2], Hà [8], Hưng [9, 10], Chơn-Hà [5, 6, 7], Minami [14], Nam [15], Hưng-Quỳnh [11], Quỳnh [18], the present author [23] and others).

It was shown that the algebraic transfer is an isomorphism for $k = 1, 2$ by Singer in [19] and for $k = 3$ by Boardman in [1]. However, for any $k \geqslant 4$, φ_k is not a monomorphism in infinitely many degrees (see Singer [19], Hưng [10].) Singer made the following conjecture.

Conjecture 1.1 (see Singer [19])**.** *The algebraic transfer φ_k is an epimorphism for any $k \geqslant 0$.*

The conjecture is true for $k \leqslant 3$. Based on the results in [22, 24], we are verifying this conjecture for $k = 4$. We hope that it is also true in this case. However, for $k \geqslant 5$, the conjecture is still open.

There is a classical operator, known as Kameko's squaring operation

$$\widetilde{Sq}^0_* : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{2d+k} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_d,$$

which is induced by an $\mathbb{F}_2$-linear map $\phi : P_k \to P_k$, given by

$$\phi(x) = \begin{cases} y, & \text{if } x = x_1x_2\ldots x_ky^2,\\ 0, & \text{otherwise,}\end{cases}$$

for any monomial $x \in P_k$. Note that ϕ is not an $\mathcal{A}$-homomorphism. However, $\phi Sq^{2i} = Sq^i\phi$ and $\phi Sq^{2i+1} = 0$ for any non-negative integer i. Since $\widetilde{Sq}^0_*$ is a homomorphism of GL_k-modules, it induces a homomorphism which is also denoted by $\widetilde{Sq}^0_* : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)^{GL_k}_{2d+k} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)^{GL_k}_d$. It was recognized by Boardman [1] for k = 3 and by Minami [14] for general k that Kameko's squaring operation commutes with the dual of the classical squaring operation on the cohomology of the Steenrod algebra, $Sq^0 : \mathrm{Ext}^{k,d+k}_{\mathcal{A}}(\mathbb{F}_2,\mathbb{F}_2) \to \mathrm{Ext}^{k,2d+2k}_{\mathcal{A}}(\mathbb{F}_2,\mathbb{F}_2)$, through the Singer algebraic transfer. This means that the following diagram is commutative:

$$\begin{array}{ccc} \mathrm{Tor}^{k,2d+2k}_{\mathcal{A}}(\mathbb{F}_2,\mathbb{F}_2) & \xrightarrow{\varphi_k} & (\mathbb{F}_2\otimes_{\mathcal{A}}P_k)^{GL_k}_{2d+k} \\ \downarrow{\scriptstyle Sq^0_*} & & \downarrow{\scriptstyle \widetilde{Sq}^0_*} \\ \mathrm{Tor}^{k,d+k}_{\mathcal{A}}(\mathbb{F}_2,\mathbb{F}_2) & \xrightarrow{\varphi_k} & (\mathbb{F}_2\otimes_{\mathcal{A}}P_k)^{GL_k}_{d}. \end{array}$$

For a positive integer n, by $\mu(n)$ one means the smallest number r for which it is possible to write $n = \sum_{1\leqslant i\leqslant r}(2^{u_i} - 1)$, where $u_i > 0$.

Theorem 1.2 (see Kameko [12])**.** *Let d be a non-negative integer. If $\mu(2d+k) = k$, then*

$$\widetilde{Sq}^0_* : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{2d+k} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_d$$

is an isomorphism of GL_k-modules.

From the result of Carlisle and Wood [3] on the boundedness conjecture, Hưng observes in [10] that, for any degree d, there exists a non-negtive integer t such that

$$(\widetilde{Sq}^0_*)^{s-t} : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{k(2^s-1)+2^sd} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{k(2^t-1)+2^td}$$

is an isomorphism of GL_k-modules for every $s \geqslant t$. However, this result does not confirm how large t should be.

Denote by $\alpha(n)$ the number of ones in dyadic expansion of a positive integer n and by $\zeta(n)$ the greatest integer u such that n is divisible by 2^u. That means $n = 2^{\zeta(n)}m$ with m an odd integer. We set

$$t(k,d) = \max\{0, k - \alpha(d+k) - \zeta(d+k)\}.$$

The following is one of our main results.

Theorem 1.3. *Let d be an arbitrary non-negative integer. Then*

$$(\widetilde{Sq}^0_*)^{s-t} : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{k(2^s-1)+2^s d} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{k(2^t-1)+2^t d}$$

is an isomorphism of GL_k-modules for every $s \geqslant t$ if and only if $t \geqslant t(k,d)$.

For either $d = 0$ or $\mu(d) \leqslant k$, we prove that $t = t(k,d)$ is the minimum number such that $\mu(k(2^s-1) + 2^s d) = k$ for every $s > t$. Then, the theorem follows from Theorem 1.2. If $\mu(d) > k$, then $\mu(k(2^s-1) + 2^s d) > k$ for every $s \geqslant 0 = t(k,d)$. From a result of Wood [28], we have $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{k(2^s-1)+2^s d} = 0$, for every $s \geqslant 0$. So, Theorem 1.3 is true for an arbitrary non-negative integer d.

It is easy to see that $t(k,d) \leqslant k-2$ for every d and $k \geqslant 2$. Hence, one gets the following.

Corollary 1.4 (See Hưng [10])**.** *Let d be an arbitrary non-negative integer. If $k \geqslant 2$, then*

$$(\widetilde{Sq}^0_*)^{s-k+2} : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{k(2^s-1)+2^s d} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{k(2^{k-2}-1)+2^{k-2} d}$$

is an isomorphism of GL_k-modules for every $s \geqslant k-2$.

Corollary 1.4 shows that the number $t = k-2$ commonly serves for every degree d. In [10], Hưng predicted that $t = k-2$ is the minimum number for this purpose and proved it for $k = 5$. It is easy to see that for $d = 2^k - k + 1$, we have $t(k,d) = k-2$. So, his prediction is true for all $k \geqslant 2$.

An application of Theorem 1.3 is the following theorem.

Theorem 1.5. *Singer's conjecture is true for $k = 5$ and the degree $5(2^s-1)$, with s an arbitrary positive integer.*

For $d = 0$, we have $t(5,0) = 3$. So, Theorem 1.3 implies that

$$(\widetilde{Sq}^0_*)^{s-3} : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{5(2^s-1)} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{35}$$

is an isomorphism of GL_5-modules for every $s \geqslant 3$. Hence, by computing the space $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)^{GL_5}_{5(2^s-1)}$ for $s = 1, 2, 3$, we obtain the following.

Theorem 1.6. *For any positive integer s, we have*

$$\dim(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)^{GL_5}_{5(2^s-1)} = \begin{cases} 0, & \text{if } s = 1, \\ 2, & \text{if } s = 2, \\ 1, & \text{if } s \geqslant 3. \end{cases}$$

This theorem has been proved in [23] for $s = 2$. In [10], Hưng also proved the theorem for $s = 2, 3$ by using a computer program of S. Shpectorov written in GAP. However, the detailed proof was unpublished at the time of the writing.

The proof Theorem 1.6 is long and very complicated. It is proved by using the admissible monomials of degree $5(2^s-1)$ in P_5. The computations are based on some results of Kameko [12] and Singer [19] on the admissible monomials and the hit monomials (see Section 2).

From the results of Chen [4], Lin [13] and Tangora [25], we have

$$\mathrm{Tor}^{\mathcal{A}}_{5,5.2^s}(\mathbb{F}_2,\mathbb{F}_2)=\begin{cases}0, & \text{if } s=1,\\ \langle (h_0^4h_4)^*, (h_1d_0)^*\rangle, & \text{if } s=2,\\ \langle (h_{s-1}d_{s-2})^*\rangle, & \text{if } s\geqslant 3,\end{cases}$$

and $h_{s-1}d_{s-2}\neq 0$, where h_{s-1} denote the Adams element in $\mathrm{Ext}^{1,2^{s-1}}_{\mathcal{A}}(\mathbb{F}_2,\mathbb{F}_2)$ and $d_{s-2}\in \mathrm{Ext}^{4,2^{s+2}+2^{s-1}}_{\mathcal{A}}(\mathbb{F}_2,\mathbb{F}_2)$ for $s\geqslant 2$. In [23], we have proved that

$$\varphi_5:\mathrm{Tor}^{\mathcal{A}}_{5,20}(\mathbb{F}_2,\mathbb{F}_2)\longrightarrow (\mathbb{F}_2\otimes_{\mathcal{A}}P_5)^{GL_5}_{15}$$

is an isomorphism. Combining the results of Hà [8] and Singer [19], we have $\varphi_5((h_{s-1}d_{s-2})^*)\neq 0$. Hence, Theorem 1.6 implies that

$$\varphi_5:\mathrm{Tor}^{\mathcal{A}}_{5,5.2^s}(\mathbb{F}_2,\mathbb{F}_2)\longrightarrow (\mathbb{F}_2\otimes_{\mathcal{A}}P_5)^{GL_5}_{5(2^s-1)}$$

is also an isomorphism for $s\geqslant 3$. Therefore, Singer's conjecture is true in the case $k=5$ and the degree $5(2^s-1)$. Theorem 1.5 is proved.

For $d=2$, we have $t(5,2)=2$ and $5(2^s-1)+2^sd=7.2^s-5$. So, by Theorem 1.3, $(\widetilde{Sq^0_*})^{s-2}:(\mathbb{F}_2\otimes_{\mathcal{A}}P_5)_{7.2^s-5}\longrightarrow(\mathbb{F}_2\otimes_{\mathcal{A}}P_5)_{23}$ is an isomorphism of GL_5-modules for every $s\geqslant 2$. Hence, by an explicit computation of $(\mathbb{F}_2\otimes_{\mathcal{A}}P_5)^{GL_5}_{7.2^s-5}$ for $s=1,2$, Tín proved in [27] the following.

Theorem 1.7 (Tín [27])**.** *Let s be a positive integer. Then, $(\mathbb{F}_2\otimes_{\mathcal{A}}P_5)^{GL_5}_{7.2^s-5}=0$.*

The theorem has been proved by Singer [19] for $s=1$. In [10], Hưng also proved this theorem for $s=2$ by using computer calculation. However, the detailed proof was also unpublished.

From the results of Tangora [25], Lin [13] and Chen [4], we can see that for any $s\geqslant 1$, $\dim\mathrm{Tor}^{\mathcal{A}}_{5,7.2^s}(\mathbb{F}_2,\mathbb{F}_2)=1$. Hence, by Theorem 1.7, the homomorphism φ_5 is an epimorphism in degree 7.2^s-5. However, it is not a monomorphism.

This paper is organized as follows. In Section 2, we recall some needed information on the admissible monomials in P_k and Singer's criterion on the hit monomials. Theorem 1.3 is proved in Section 3. In Section 4, we explicitly determine a system of $\mathcal{A}$-generators for P_5 in degree $5(2^s-1)$. Theorem 1.6 is proved in Section 5 by using the results in Section 4. Finally, in the appendix we list the admissible monomials of degrees 5, 7, 15, 16, 35 in P_4 and P_5.

Theorems 1.3 and 1.7 have already been announced in [26].

2. Preliminaries

In this section, we recall some needed information from Kameko [12] and Singer [20], which will be used in the next sections.

Notation 2.1. Let $\alpha_i(a)$ denote the i-th coefficient in dyadic expansion of a non-negative integer a. That means $a=\alpha_0(a)2^0+\alpha_1(a)2^1+\alpha_2(a)2^2+\ldots$, for $\alpha_i(a)=0$ or 1 with $i\geqslant 0$.

For a set of integers $\mathbb{J} = \{j_1, j_2, \ldots, j_s\}$ with $1 \leqslant j_u \leqslant k$, $1 \leqslant u \leqslant s$, we define the monomial $X_{\mathbb{J}} \in P_k$ by setting $X_{\mathbb{J}} = \prod_{j\neq j_u, \forall u} x_j = x_1 \ldots \hat{x}_{j_1} \ldots \hat{x}_{j_s} \ldots x_k$.

Let $x = x_1^{a_1} x_2^{a_2} \ldots x_k^{a_k} \in P_k$. Denote $\nu_j(x) = a_j, 1 \leqslant j \leqslant k$. Set

$$\mathbb{J}_t(x) = \{j : 1 \leqslant j \leqslant k,\ \alpha_t(\nu_j(x)) = 0\},$$

for $t \geqslant 0$. Then, we have $x = \prod_{t\geqslant 0} X_{\mathbb{J}_t(x)}^{2^t}$.

Definition 2.2. For a monomial x in P_k, define two sequences associated with x by

$$\omega(x) = (\omega_1(x), \omega_2(x), \ldots, \omega_i(x), \ldots), \quad \sigma(x) = (\nu_1(x), \nu_2(x), \ldots, \nu_k(x)),$$

where $\omega_i(x) = \sum_{1\leqslant j\leqslant k} \alpha_{i-1}(\nu_j(x)) = \deg X_{\mathbb{J}_{i-1}(x)},\ i \geqslant 1$. The sequence $\omega(x)$ is called the weight vector of x.

The sequence $\omega = (\omega_1, \omega_2, \ldots, \omega_i, \ldots)$ of non-negative integers is called the weight vector if $\omega_i = 0$ for $i \gg 0$.

The sets of the weight vectors and the exponent vectors are given the left lexicographical order.

For a weight vector ω, we define $\deg \omega = \sum_{i>0} 2^{i-1}\omega_i$. Denote by $P_k(\omega)$ the subspace of P_k spanned by all monomials y such that $\deg y = \deg \omega$, $\omega(y) \leqslant \omega$, and by $P_k^-(\omega)$ the subspace of P_k spanned by all monomials $y \in P_k(\omega)$ such that $\omega(y) < \omega$.

Definition 2.3. Let ω be a weight vector and f, g two polynomials of the same degree in P_k.

i) $f \equiv g$ if and only if $f + g \in \mathcal{A}^+P_k$. If $f \equiv 0$ then f is called *hit*.

ii) $f \equiv_\omega g$ if and only if $f + g \in \mathcal{A}^+P_k + P_k^-(\omega)$.

Obviously, the relations $\equiv$ and $\equiv_\omega$ are equivalence ones. Denote by $QP_k(\omega)$ the quotient of $P_k(\omega)$ by the equivalence relation $\equiv_\omega$. Then, we have

$$QP_k(\omega) = P_k(\omega)/((\mathcal{A}^+P_k \cap P_k(\omega)) + P_k^-(\omega)).$$

For a polynomial $f \in P_k$, we denote by $[f]$ the class in $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k = P_k/\mathcal{A}^+P_k$ represented by f. If ω is a weight vector, then denote by $[f]_\omega$ the class in the space $P_k/(\mathcal{A}^+P_k + P_k^-(\omega))$ represented by f .

Denote by $|S|$ the cardinal of a set S. If S is a subset of a vector space, then we denote by $\langle S\rangle$ the subspace spanned by S.

It is easy to see that

$$QP_k(\omega) \cong QP_k^\omega := \langle\{[x] \in QP_k : x \text{ is admissible and } \omega(x) = \omega\}\rangle.$$

So, we get

$$(\mathbb{F}_2\otimes_{\mathcal{A}}P_k)_n = \bigoplus_{\deg\omega=n} QP_k^\omega \cong \bigoplus_{\deg\omega=n} QP_k(\omega).$$

Hence, we can identify the vector space $QP_k(\omega)$ with $QP_k^\omega \subset QP_k$.

We note that the weight vector of a monomial is invariant under the permutation of the generators x_i, hence $QP_k(\omega)$ is an Σ_k-module, where $\Sigma_k \subset GL_k$ is the symmetric group. Furthermore, we have the following.

Lemma 2.4. *Let ω be a weight vector. Then, $QP_k(\omega)$ is an GL_k-module.*

Proof. We prove the lemma by proving the fact that if x is a monomial in P_k, then $g_k(x) \in P_k(\omega(x))$.

If $\nu_1(x) = 0$, then $x = g_k(x)$ and $\omega(g_k(x)) = \omega(x)$. Suppose $\nu_1(x) > 0$ and $\nu_1(x) = 2^{t_1} + \ldots + 2^{t_b}$, where $0 \leqslant t_1 < \ldots < t_b,\ b \geqslant 1$.

Since $x = \prod_{t\geqslant 0} X_{\mathbb{J}_t(x)}^{2^t} \in P_k$ and g_k is a homomorphism of algebras, we have

$$g_k(x) = \prod_{t\geqslant 0}(g_k(X_{\mathbb{J}_t(x)}))^{2^t} = \left(\prod_{u=1}^{b}\left((x_1+x_2)X_{\mathbb{J}_{t_u}(x)\cup 1}\right)^{2^{t_u}}\right)\left(\prod_{t\neq t_1,t_2,\ldots,t_b} X_{\mathbb{J}_t(x)}^{2^t}\right).$$

Then, $g_k(x)$ is a sum of monomials of the form

$$\bar{y} = \left(\prod_{j=1}^{c}\left(x_2 X_{\mathbb{J}_{t_{u_j}}(x)\cup 1}\right)^{2^{t_u}}\right)\left(\prod_{t\neq t_{u_1},\ldots,t_{u_c}} X_{\mathbb{J}_t(x)}^{2^t}\right),$$

where $0 \leqslant c \leqslant b$. If $c = 0$, then $\bar{y} = x$ and $\omega(\bar{y}) = \omega(x)$. Suppose $c > 0$.

If $2 \in \mathbb{J}_{t_{u_j}}(x)$ for all j, $1 \leqslant j \leqslant c$, then $\omega(\bar{y}) = \omega(x)$ and $\bar{y} \in P_k(\omega(x))$. Suppose there is an index j such that $2 \notin \mathbb{J}_{t_{u_j}}(x)$. Let j_0 be the smallest index such that $2 \notin \mathbb{J}_{t_{u_{j_0}}}(x)$. Then, we have

$$\omega_i(\bar{y}) = \begin{cases} \omega_i(x), & \text{if } i \leqslant t_{u_{j_0}}, \\ \omega_i(x) - 2, & \text{if } i = t_{u_{j_0}} + 1. \end{cases}$$

Hence $\omega(\bar{y}) < \omega(x)$ and $\bar{y} \in P_k(\omega(x))$. The lemma is proved. □

Note that $V_k \cong \langle x_1, x_2, \ldots, x_k\rangle \subset P_k$. For $1 \leqslant i \leqslant k$, define the $\mathbb{F}_2$-linear map $g_i : V_k \to V_k$, which is determined by $g_i(x_i) = x_{i+1}, g_i(x_{i+1}) = x_i,\ g_i(x_j) = x_j$ for $j \neq i, i+1,\ 1 \leqslant i < k$, and $g_k(x_1) = x_1 + x_2,\ g_k(x_j) = x_j$ for $j > 1$. The general linear group $GL_k \cong GL(V_k)$ is generated by $g_i,\ 1 \leqslant i \leqslant k$, and the symmetric group Σ_k is generated by $g_i,\ 1 \leqslant i < k$. The $\mathbb{F}_2$-linear map g_i induces a homomorphism of $\mathcal{A}$-algebras which is also denoted by $g_i : P_k \to P_k$. So, an element $[f]_\omega \in QP_k(\omega)$ is an GL_k-invariant if and only if $g_i(f) \equiv_\omega f$ for $1 \leqslant i \leqslant k$. It is an Σ_k-invariant if and only if $g_i(f) \equiv_\omega f$ for $1 \leqslant i < k$.

Definition 2.5. Let x, y be monomials of the same degree in P_k. We say that $x < y$ if and only if one of the following holds:

i) $\omega(x) < \omega(y)$;

ii) $\omega(x) = \omega(y)$ and $\sigma(x) < \sigma(y)$.

Definition 2.6. A monomial x is said to be inadmissible if there exist monomials $y_1, y_2, \ldots, y_t$ such that $y_j < x$ for $j = 1, 2, \ldots, t$ and $x + \sum_{j=1}^{t} y_j \in \mathcal{A}^+P_k$. A monomial x is said to be admissible if it is not inadmissible.

Obviously, the set of all the admissible monomials of degree n in P_k is a minimal set of $\mathcal{A}$-generators for P_k in degree n.

Definition 2.7. A monomial x in P_k is said to be strictly inadmissible if and only if there exist monomials $y_1, y_2, \ldots, y_t$ such that $y_j < x$, for $j = 1, 2, \ldots, t$ and

$$x = \sum_{j=1}^{t} y_j + \sum_{u=1}^{2^s-1} Sq^u(q_u)$$

with $s = \max\{i : \omega_i(x) > 0\}$ and suitable polynomials $q_u \in P_k$.

It is easy to see that if x is strictly inadmissible, then it is inadmissible.

Theorem 2.8 (See Kameko [12]). *Let x, y, w be monomials in P_k such that $\omega_i(x) = 0$ for $i > r > 0$, $\omega_s(w) \neq 0$ and $\omega_i(w) = 0$ for $i > s > 0$.*

i) *If w is inadmissible, then xw^{2^r} is also inadmissible.*

ii) *If w is strictly inadmissible, then wy^{2^s} is also strictly inadmissible.*

Now, we recall a result of Singer [20] on the hit monomials in P_k.

Definition 2.9. A monomial z in P_k is called a spike if $\nu_j(z) = 2^{t_j} - 1$ for t_j a non-negative integer and $j = 1, 2, \ldots, k$. If z is a spike with $t_1 > t_2 > \ldots > t_{r-1} \geqslant t_r > 0$ and $t_j = 0$ for $j > r$, then it is called the minimal spike.

In [20], Singer showed that if $\mu(n) \leqslant k$, then there exists uniquely a minimal spike of degree n in P_k.

Lemma 2.10 (See [17]). *All the spikes in P_k are admissible and their weight vectors are weakly decreasing. Furthermore, if a weight vector ω is weakly decreasing and $\omega_1 \leqslant k$, then there is a spike z in P_k such that $\omega(z) = \omega$.*

The following is a criterion for the hit monomials in P_k.

Theorem 2.11 (See Singer [20]). *Suppose $x \in P_k$ is a monomial of degree n, where $\mu(n) \leqslant k$. Let z be the minimal spike of degree n. If $\omega(x) < \omega(z)$, then x is hit.*

This result implies the one of Wood, which originally is a conjecture of Peterson [16].

Theorem 2.12 (See Wood [28]). *If $\mu(n) > k$, then $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_n = 0$.*

Now, we recall some notations and definitions in [24], which will be used in the next sections. We set

$$P_k^0 = \langle \{x = x_1^{a_1} x_2^{a_2} \ldots x_k^{a_k} \ : \ a_1 a_2 \ldots a_k = 0\} \rangle,$$
$$P_k^+ = \langle \{x = x_1^{a_1} x_2^{a_2} \ldots x_k^{a_k} \ : \ a_1 a_2 \ldots a_k > 0\} \rangle.$$

It is easy to see that P_k^0 and P_k^+ are the $\mathcal{A}$-submodules of P_k. Furthermore, we have the following.

Proposition 2.13. *We have a direct summand decomposition of the $\mathbb{F}_2$-vector spaces $\mathbb{F}_2 \otimes_{\mathcal{A}} P_k = QP_k^0 \oplus QP_k^+$. Here $QP_k^0 = \mathbb{F}_2 \otimes_{\mathcal{A}} P_k^0$ and $QP_k^+ = \mathbb{F}_2 \otimes_{\mathcal{A}} P_k^+$.*

Definition 2.14. For $1 \leqslant i \leqslant k$, define the homomorphism $f_i : P_{k-1} \to P_k$ of algebras by substituting

$$f_i(x_j) = \begin{cases} x_j, & \text{if } 1 \leqslant j < i, \\ x_{j+1}, & \text{if } i \leqslant j < k. \end{cases}$$

Obviously, we have the following.

Proposition 2.15. *If B is a minimal set of generators for $\mathcal{A}$-module P_{k-1} in degree n, then $f(B) := \bigcup_{1 \leqslant i \leqslant k} f_i(B)$ is also a minimal set of generators for $\mathcal{A}$-module P_k^0 in degree n.*

Definition 2.16. For any $1 \leqslant i < j \leqslant k$, we define the homomorphism $p_{(i;j)} : P_k \to P_{k-1}$ of algebras by substituting

$$p_{(i;j)}(x_u) = \begin{cases} x_u, & \text{if } 1 \leqslant u < i, \\ x_{j-1}, & \text{if } u = i, \\ x_{u-1}, & \text{if } i < u \leqslant k. \end{cases}$$

Then, $p_{(i;j)}$ is a homomorphism of $\mathcal{A}$-modules. In particular, $p_{(i;j)}(f_i(y)) = y$ for any $y \in P_{k-1}$.

Lemma 2.17 (see [17])**.** *Let x be a monomial in P_k. Then, $p_{(i;j)}(x) \in P_{k-1}(\omega(x))$.*

Lemma 2.17 implies that if ω is a weight vector and $x \in P_k(\omega)$, then $p_{(i;j)}(x) \in P_{k-1}(\omega)$. Moreover, $p_{(i;j)}$ passes to a homomorphism from $QP_k(\omega)$ to $QP_{k-1}(\omega)$.

For a subset $B \subset P_k$ and a weight vector ω, we denote $[B] = \{[f] : f \in B\}$ and $[B]_\omega = \{[f]_\omega : f \in B\}$. From Theorem 2.11, we see that if ω is the weight vector of a minimal spike in P_k, then $[B]_\omega = [B]$.

From now on, we denote by $B_k(n)$ the set of all admissible monomials of degree n in P_k, $B_k^0(n) = B_k(n) \cap P_k^0$, $B_k^+(n) = B_k(n) \cap P_k^+$. For a weight vector ω of degree n, we set $B_k(\omega) = B_k(n) \cap P_k(\omega)$, $B_k^+(\omega) = B_k^+(n) \cap P_k(\omega)$. Then, $[B_k(\omega)]_\omega$ and $[B_k^+(\omega)]_\omega$, are respectively the basses of the $\mathbb{F}_2$-vector spaces $QP_k(\omega)$ and $QP_k^+(\omega) := QP_k(\omega) \cap QP_k^+$.

For any monomials $z, z_1, z_2, \ldots, z_m$ in $P_k(\omega)$ with $m \geqslant 1$, and for a subgroup $G \subset GL_k$, we denote

$$\begin{aligned} G(z_1, z_2, \ldots, z_m) &= \{\sigma z_t : \sigma \in G, 1 \leqslant t \leqslant m\} \subset P_k(\omega), \\ [B(z_1, z_2, \ldots, z_m)]_\omega &= [B_k(\omega)]_\omega \cap \langle [\Sigma_k(z_1, z_2, \ldots, z_m)]_\omega \rangle, \\ p(z) &= \sum_{y \in B_k(n) \cap \Sigma_k(z)} y. \end{aligned}$$

We note that the $\mathbb{F}_2$-vector subspace $\langle [G(z_1, z_2, \ldots, z_m)]_\omega \rangle$ is the G-submodule of $QP_k(\omega)$ generated by the set $\{[z_1]_\omega, [z_2]_\omega, \ldots, [z_m]_\omega\}$.

3. Proof of Theorem 1.3

To make the paper self-contained, we give here a proof for the following lemma, which is an elementary property of the μ-function.

Lemma 3.1. *Let n be a positive integer. Then, $\mu(n) = s$ if and only if there exists uniquely a sequence of integers $v_1 > v_2 > \ldots > v_{s-1} \geqslant v_s > 0$ such that*

$$n = 2^{v_1} + 2^{v_2} + \ldots + 2^{v_{s-1}} + 2^{v_s} - s = \sum_{i=1}^{s} (2^{v_i} - 1). \tag{3.1}$$

Proof. Assume that $\mu(n) = s$. Set $\beta(n) = \min\{u \in \mathbb{N} : \alpha(n+u) \leqslant u\}$. We prove $\mu(n) = \beta(n)$.

Suppose $\beta(n) = t$. Then $\alpha(n+t) = r \leqslant t$, and $n = 2^{c_1} + 2^{c_2} + \ldots + 2^{c_r} - t$, where $c_1 > c_2 > \ldots > c_r \geqslant 0$.

If $c_r \leqslant t - r$ then

$$\alpha(n + t - 1) = \alpha(2^{c_1} + 2^{c_2} + \ldots + 2^{c_{r-1}} + 2^{c_r} - 1) = r - 1 + c_r \leqslant t - 1.$$

Hence $\beta(n) \leqslant t - 1$. This contradicts the fact that $\beta(n) = t$. So, $c_r > t - r$.

If $r = t$ then $c_t = c_r > t - r = 0$. Set $v_i = c_i,\ i = 1, 2, \ldots, t$. We obtain

$$n = 2^{v_1} + 2^{v_2} + \ldots + 2^{v_{t-1}} + 2^{v_t} - t = \sum_{i=1}^{t}(2^{v_i} - 1),$$

where $v_1 > v_2 > \ldots > v_{t-2} > v_{t-1} > v_t > 0$. Hence, $\mu(n) \leqslant t = \beta(n)$.

Suppose $r < t$. Obviously,

$$2^{c_r} = 2^{c_r - 1} + \ldots + 2^{c_r - t + r + 1} + 2^{c_r - t + r} + 2^{c_r - t + r}.$$

Set

$$\begin{aligned} &v_i = c_i,\ i = 1, 2, \ldots, r - 1,\\ &v_{r+\ell} = c_r - \ell - 1 > 0,\ \ell = 0, 1, \ldots, t - r - 2,\\ &v_{t-1} = v_t = c_r - t + r > 0. \end{aligned}$$

Then, we get

$$n = 2^{v_1} + 2^{v_2} + \ldots + 2^{v_{t-1}} + 2^{v_t} - t = \sum_{i=1}^{t}(2^{v_i} - 1),$$

with $v_1 > v_2 > \ldots > v_{t-2} > v_{t-1} = v_t > 0$. Hence $\mu(n) \leqslant t = \beta(n)$.

Since $\mu(n) = s$, $n = \sum_{i=1}^{s}(2^{h_i} - 1)$ with h_i positive integers. Then, $\alpha(n + s) = \alpha(\sum_{i=1}^{s} 2^{h_i}) \leqslant s$. So, we get $\mu(n) = s \geqslant \beta(n)$. Hence, $t = \beta(n) = \mu(n) = s$. Thus, n is of the form (3.1).

Now, assume that n is of the form (3.1). Then, $\mu(n) \leqslant s$. We prove $\mu(n) = s$ by induction on s.

If $s = 1$, then $\mu(n) = 1$, since $\mu(n) > 0$. If $s = 2$, then $\alpha(n+1) = \alpha(2^{v_1} + 2^{v_2} - 1) = 1 + v_2 > 1$. Hence, $\mu(n) = \beta(n) \geqslant 2$. So, $\mu(n) = 2$.

Suppose $s > 2$. By the inductive hypothesis, $\mu(n + 1 - 2^{v_1}) = s - 1$.

It is well-known that there exists uniquely an integer d such that $2^d \leqslant n + 1 < 2^{d+1}$. Since $v_1 > v_2 > \ldots > v_{s-1} \geqslant v_s > 0$, we have

$$2^{v_1} \leqslant n + 1 < 2^{v_1} + 2^{v_1 - 1} + \ldots + 2^{v_1 - s + 2} + 2^{v_1 - s + 2} = 2^{v_1 + 1}.$$

So, we get $v_1 = d$. Set $\mu(n) = t \leqslant s$. There exists $u_1 > u_2 > \ldots > u_{t-1} \geqslant u_t > 0$ such that $n = 2^{u_1} + 2^{u_2} + \ldots + 2^{u_t} - t$. Then, $u_1 = d = v_1$ and $\alpha(n + 1 - 2^d + t - 1) \leqslant t - 1$. Hence, $t - 1 \geqslant \beta(n + 1 - 2^d) = \mu(n + 1 - 2^d) = s - 1$. This implies $t \geqslant s$ and $\mu(n) = t = s$.

By induction on i, we get $u_i = v_i$ for $1 \leqslant i \leqslant s$. The lemma is proved. □

From this lemma we easily obtain the following.

Corollary 3.2 (See Kameko [12]). *Let n, k be positive integers. Then*

i) *$\mu(n) > k$ if and only if $\alpha(n + k) > k$.*

ii) *If $n > \mu(n)$, then $n - \mu(n)$ is even and $\mu\big(\frac{n-\mu(n)}{2}\big) \leqslant \mu(n)$.*

iii) *$\mu\big(2n + \mu(n)\big) = \mu(n)$.*

Suppose that d is a non-negative integer such that $\mu(2d + k) = s < k$. By Lemma 3.1, there exists a sequence of integers $v_1 > v_2 > \ldots > v_{s-1} \geqslant v_s > 0$ such that $2d + k = \sum_{i=1}^{s}(2^{v_i} - 1)$. Set $z = x_1^{2^{v_1}-1} x_2^{2^{v_2}-1} \ldots x_s^{2^{v_s}-1} \in (P_k)_{2d+k}$. Since z is a spike and $s < k$, we have $[z] \neq 0$ and $\widetilde{Sq}^0_*([z]) = 0$. So, one gets the following.

Corollary 3.3. *Let d be an arbitrary non-negative integer. If $\mu(2d+k) < k$, then*

$$(\widetilde{Sq}^0_*)_{(k,d)} := \widetilde{Sq}^0_* : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{2d+k} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_d$$

is not a monomorphism.

Now, we are ready to prove Theorem 1.3.

Proof of Theorem 1.3. Set $q = \alpha(d+k)$, $r = \zeta(d+k)$ and $m = k(2^t-1) + 2^t d$. From the proof of Lemma 3.1 and Corollary 3.2, we see that if $q > k$, then

$$\mu(k(2^s-1) + 2^s d) \geqslant \mu(d) > k$$

for any $s \geqslant 0 = t(k,d)$. By Theorem 2.12,

$$(\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{k(2^s-1)+2^s d} = 0,\ (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_d = 0.$$

So, the theorem holds.

Assume that $q \leqslant k$. Using Theorem 1.2, Corollaries 3.2 and 3.3, we see that the homomorphism

$$(\widetilde{Sq}^0_*)^{s-t} : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{k(2^s-1)+2^s d} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{k(2^t-1)+2^t d}$$

is an isomorphism of GL_k-modules for every $s \geqslant t$ if and only if $\mu(2m+k) = k$.

Since $\alpha(d+k) = q$ and $\zeta(d+k) = r$, there exists a sequence of integers $c_1 > c_2 > \ldots > c_{q-1} > c_q = r \geqslant 0$ such that

$$d + k = 2^{c_1} + 2^{c_2} + \ldots + 2^{c_q}.$$

If $q = k$, then

$$\begin{aligned} 2m + k &= k(2^{t+1}-1) + 2^{t+1}d = k(2^{t+1}-1) + 2^{t+1}(2^{c_1} + 2^{c_2} + \ldots + 2^{c_k} - k) \\ &= 2^{c_1+t+1} + 2^{c_2+t+1} + \ldots + 2^{c_k+t+1} - k. \end{aligned}$$

By Lemma 3.1, $\mu(2m+k) = k$ for any $t \geqslant 0 = t(k,d)$. Hence, the theorem holds.

Suppose that $q < k$. Then, we have

$$\begin{aligned} 2m + k &= 2^{c_1+t+1} + 2^{c_2+t+1} + \ldots + 2^{c_{q-1}+t+1} + 2^{r+t+1} - k \\ &= 2^{c_1+t+1} + 2^{c_2+t+1} + \ldots + 2^{c_{q-1}+t+1} \\ &\quad + 2^{r+t} + 2^{r+t-1} + \ldots + 2^{r+t-(k-q-1)} + 2^{r+t-(k-q-1)} - k. \end{aligned} \tag{3.2}$$

If $q + r \geqslant k$, then $r + t - (k-q-1) = q + r - k + 1 + t > 0$ for any $t \geqslant 0 = t(k,d)$. By Lemma 3.1, $\mu(2m+k) = k$. So, the theorem is true.

If $q + r < k$, then from Lemma 3.1 and the relation (3.2), we see that $\mu(2m+k) = k$ if and only if $t \geqslant k - q - r = t(k,d)$. The theorem is completely proved. □

4. $\mathcal{A}$-generators for P_5 in degree $5(2^s-1)$

To prove Theorem 1.6, we need to determine a system of $\mathcal{A}$-generators for P_5 in degree $5(2^s-1)$. From Theorem 1.3, we see that

$$(\widetilde{Sq}^0_*)^{s-3} : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{5(2^s-1)} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{35}$$

is an isomorphism of GL_5-modules for every $s \geqslant 3$. So, we need only to determine $\mathcal{A}$-generators for P_5 in degree $5(2^s-1)$ for $s = 1, 2, 3$.

4.1. **The Cases $s = 1, 2$.**

By using a result in [24], we can easily obtain the following.

Proposition 4.1.1. *There exist exactly* 46 *admissible monomials of degree* 5 *in* P_5. *Consequently* $\dim(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_5 = 46$.

For $s = 2$, we have $5(2^s - 1) = 15$. The space $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{15}$ has been computed in [23].

Proposition 4.1.2 (See [23])**.** *There exist exactly* 432 *admissible monomials of degree* 15 *in* P_5. *Consequently* $\dim(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{15} = 432$.

The admissible monomials of degrees 5 and 15 are explicitly determined as in Subsections 6.1 and 6.2.

4.2. **The admissible monomials of degree** 16 **in** P_5**.**

To compute $\mathcal{A}$-generators for P_5 in degree $5(2^s - 1)$ for $s = 3$, we need to determine the admissible monomials of degree 16 in P_5.

Lemma 4.2.1. *If x is an admissible monomial of degree* 16 *in P_5, then $\omega(x)$ is one of the following sequences:*

$$(2, 1, 1, 1),\ (2, 1, 3),\ (2, 3, 2),\ (4, 2, 2),\ (4, 4, 1).$$

Proof. Observe that $z = x_1^{15}x_2$ is the minimal spike of degree 16 in P_5 and $\omega(z) = (2, 1, 1, 1)$. Since $[x] \neq 0$, by Theorem 2.11, either $\omega_1(x) = 2$ or $\omega_1(x) = 2$. If $\omega_1(x) = 2$, then $x = x_ix_jy^2$ with y a monomial of degree 7 in P_5. Since x is admissible, by Theorem 2.8, y is admissible. A routine computation shows that either $\omega(y) = (1, 1, 1)$ or $\omega(y) = (1, 3)$ or $\omega(y) = (3, 2)$. If $\omega_1(x) = 4$, then $x = X_jy_1^2$ with y_1 an admissible monomial of degree 6 in P_5. It is easy to see that either $\omega(y_1) = (2, 2)$ or $\omega(y_1) = (4, 1)$. The lemma is proved. □

We have $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{16} = (QP_5^0)_{16} \bigoplus (QP_5^+)_{16}$. By Lemma 4.2.1,

$$\begin{aligned}(QP_5^+)_{16} \cong\ & QP_5^+(2, 1, 1, 1) \bigoplus QP_5^+(2, 1, 3)\\ & \bigoplus QP_5^+(2, 3, 2) \bigoplus QP_5^+(4, 2, 2) \bigoplus QP_5^+(4, 4, 1).\end{aligned}$$

From a result in [24], we easily obtain $\dim(QP_5^0)_{16} = 255$.

Proposition 4.2.2. *$(QP_5^+)_{16}$ is the $\mathbb{F}_2$-vector space of dimension* 188 *with a basis consisting of all the classes represented by the monomials $a_t = a_{16,t},\ 1 \leqslant t \leqslant 188$, which are determined as in Subsection 6.4.*

To prove this proposition, we need the following lemma which is easily proved by a direct computation.

Lemma 4.2.3. *The following monomials are strictly inadmissible:*

i) $x_j^2x_\ell x_t,\ j < \ell < t$; $x_j^2x_\ell x_t^2x_u^3,\ j < \ell$; $x_j^2x_\ell x_tx_u^2x_v^2,\ j < \ell$; $x_j^3x_\ell^{12}x_t,\ x_j^3x_\ell^4x_t^9,\ x_j^3x_\ell^5x_t^8,\ x_j^3x_\ell^4x_tx_u^8,\ j < \ell < t$; $x_j^2x_\ell^3x_t^3$.

ii) $x_j^3x_\ell^4x_t^4x_u^5,\ x_j^3x_\ell^4x_tx_u^4x_v^4,\ j < \ell < t$.

iii) $x_jx_\ell^6x_t^3x_u^6,\ x_jx_\ell^6x_tx_u^2x_v^6,\ j < \ell < t$; $x_jx_\ell^2x_t^2x_u^5x_v^6, u \geqslant 4$; $x_jx_\ell^2x_t^3x_u^4x_v^6, t \geqslant 3$; $x_jx_\ell^2x_t^6x_u^7,\ x_jx_\ell^2x_t^2x_u^4x_v^7$; $x_1^3x_2^4x_3x_4^2x_5^6,\ x_1^3x_2^4x_3x_4^6x_5^2$.

iv) $x_j^2x_\ell x_t^3x_u^3x_v^3,\ j < \ell$; $x_j^2x_\ell x_tx_ux_v^3,\ j < \ell < t < u$; $x_1^3x_2^4x_3^3x_4^3x_5^3$.

Here (j, ℓ, t, u, v) is a permutation of $(1, 2, 3, 4, 5)$.

Proof of Proposition 4.2.2. From Lemma 4.2.1, we have

$$B_5^+(16) = B_5^+(2,1,1,1) \cup B_5^+(2,1,3) \cup B_5^+(2,3,2) \cup B_5^+(4,2,2) \cup B_5^+(4,4,1).$$

We prove $|B_5^+(2,1,1,1)| = 4$, $|B_5^+(2,1,3)| = 5$, $|B_5^+(2,3,2)| = 20$, $|B_5^+(4,2,2)| = 110$ and $|B_5^+(4,4,1)| = 49$. For simplicity, we prove $|B_5^+(2,3,2)| = 20$ by showing that $B_5^+(2,3,2) = \{a_t = a_{16,t} : 10 \leqslant t \leqslant 29\}$. The others are proved by the similar computations.

Let x be an admissible monomial in P_5^+ such that $\omega(x) = (2,3,2) := \omega$. Then $x = x_j x_\ell y^2$ with $1 \leqslant j < \ell \leqslant 5$ and y a monomial of degree 7 in P_5. Since x is admissible, by Theorem 2.8, $y \in B_5(3,2)$.

Let $z \in B_5(3,2)$ such that $x_j x_\ell z^2 \in P_5^+$. By a direct computation, we see that if $x_j x_\ell z^2 \neq a_t$, for all $t,\ 10 \leqslant t \leqslant 29$, then there is a monomial w which is given in Lemma 4.2.3 such that $x_j x_\ell z^2 = w z_1^{2^u}$ with suitable monomial $z_1 \in P_5$, and $u = \max\{j \in \mathbb{Z} : \omega_j(w) > 0\}$. By Theorem 2.8, $x_j x_\ell z^2$ is inadmissible. Since $x = x_j x_\ell y^2$ with $y \in B_5(3,2)$ and x is admissible, one obtain $x = a_t$ for some $t,\ 10 \leqslant t \leqslant 29$. Hence, $QP_5^+(\omega)$ is spanned by the set $\{[a_t] : 10 \leqslant t \leqslant 29\}$.

We now prove the set $\{[a_t] : 10 \leqslant t \leqslant 29\}$ is linearly independent in $QP_5(\omega)$. Suppose there is a linear relation $\mathcal{S} = \sum_{t=10}^{29} \gamma_t a_t \equiv_\omega 0$, where $\gamma_t \in \mathbb{F}_2$.

From a result in [24], $\dim QP_4^+(2,3,2) = 4$, with the basis $\{[w_u] : 1 \leqslant u \leqslant 4\}$, where

$$w_1 = x_1x_2^3x_3^6x_4^6,\ w_2 = x_1^3x_2x_3^6x_4^6,\ w_3 = x_1^3x_2^5x_3^2x_4^6,\ w_4 = x_1^3x_2^5x_3^6x_4^2.$$

By a direct computation using Lemma 2.17, we get

$$\begin{aligned} p_{(1;2)}(\mathcal{S}) &\equiv_\omega \gamma_{13}w_2 + \gamma_{14}w_3 + \gamma_{15}w_4 \equiv_\omega 0,\\ p_{(1;3)}(\mathcal{S}) &\equiv_\omega \gamma_{10}w_1 + \gamma_{18}w_3 + \gamma_{19}w_4 \equiv_\omega 0. \end{aligned}$$

The above relations imply $\gamma_t = 0$ for $t = 10, 13, 14, 15, 18, 19$. Then,

$$\begin{aligned} p_{(1;4)}(\mathcal{S}) &\equiv_\omega \gamma_{11}w_1 + \gamma_{16}w_2 + \gamma_{21}w_4 \equiv_\omega 0,\\ p_{(1;5)}(\mathcal{S}) &\equiv_\omega \gamma_{12}w_1 + \gamma_{17}w_2 + \gamma_{20}w_3 \equiv_\omega 0,\\ p_{(2;3)}(\mathcal{S}) &\equiv_\omega \gamma_{24}w_3 + \gamma_{25}w_4 \equiv_\omega 0,\\ p_{(2;4)}(\mathcal{S}) &\equiv_\omega \gamma_{11}w_1 + \gamma_{22}w_2 + \gamma_{27}w_4 \equiv_\omega 0,\\ p_{(2;5)}(\mathcal{S}) &\equiv_\omega \gamma_{12}w_1 + \gamma_{23}w_2 + \gamma_{26}w_3 \equiv_\omega 0. \end{aligned}$$

From the last equalities, we obtain $\gamma_t = 0$ for $t \neq 28, 29$. Then,

$$p_{(3;4)}(\mathcal{S}) \equiv_\omega \gamma_{29}w_4 \equiv_\omega 0,\ p_{(3;5)}(\mathcal{S}) \equiv_\omega \gamma_{28}w_3 \equiv_\omega 0.$$

So, $\gamma_t = 0$ for all $t,\ 10 \leqslant t \leqslant 29$, completing the proof. □

By combining the above results, one gets the following.

Corollary 4.2.4. *There exist exactly* 443 *admissible monomials of degree* 16 *in* P_5. *Consequently* $\dim(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{16} = 443$.

4.3. **The Case $s = 3$.**

For $s = 3$, we have $5(2^s - 1) = 35$. Since Kameko's squaring operation

$$(\widetilde{Sq^0_*})_{(5,15)} : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{35} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{15}$$

is an epimorphism, we have $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{35} \cong \operatorname{Ker}(\widetilde{Sq^0_*})_{(5,15)} \bigoplus (\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{15}$. Hence, we need only to compute $\operatorname{Ker}(\widetilde{Sq^0_*})_{(5,15)}$.

Lemma 4.3.1. *If x is an admissible monomial of degree 35 in P_5 and $[x] \in \mathrm{Ker}(\widetilde{Sq}^0_*)_{(5,15)}$, then $\omega(x)$ is one of the following sequences:*

$$\omega_{(1)} = (3,2,1,1,1),\ \omega_{(2)} = (3,2,1,3),\ \omega_{(3)} = (3,2,3,2),$$
$$\omega_{(4)} = (3,4,2,2),\ \omega_{(5)} = (3,4,4,1).$$

Proof. Note that $z = x_1^{31}x_2^3x_3$ is the minimal spike of degree 35 in P_5 and $\omega(z) = (3,2,1,1,1)$. Since $[x] \neq 0$, by Theorem 2.11, either $\omega_1(x) = 3$ or $\omega_1(x) = 5$. If $\omega_1(x) = 5$, then $x = X_\emptyset y^2$ with y a monomial of degree 15 in P_5. Since x is admissible, by Theorem 2.8, y is admissible. Hence, $(\widetilde{Sq}^0_*)_{(5,15)}([x]) = [y] \neq 0$. This contradicts the fact that $[x] \in \mathrm{Ker}(\widetilde{Sq}^0_*)_{(5,15)}$, so $\omega_1(x) = 3$. Then, we have $x = x_ix_jx_\ell y_1^2$ with y_1 an admissible monomial of degree 16 in P_5. Now, the lemma follows from Lemma 4.2.1. □

From Lemma 4.3.1 and a result in [24], we obtain

$$\begin{aligned}
&\mathrm{Ker}(\widetilde{Sq}^0_*)_{(5,15)} = \bigoplus_{j=1}^{5} QP_5(\omega_{(j)}),\\
&QP_5(\omega_{(1)}) = (QP_5^0)_{35} \bigoplus QP_5^+(\omega_{(1)}),\\
&QP_5(\omega_{(j)}) = QP_5^+(\omega_{(j)}),\ j = 2,3,4,5,\\
&\dim(QP_5^0)_{35} = 460.
\end{aligned}$$

Proposition 4.3.2. *There exist exactly 160 admissible monomials in P_5^+ such that their weight vectors are $\omega_{(1)}$. Consequently $\dim QP_5^+(\omega_{(1)}) = 160$.*

We denote the monomials in $B_5(\omega_{(1)})$ by $a_t = a_{35,t},\ 1 \leqslant t \leqslant 160$, as given in Subsection 6.5. We need some lemmas for the proof of this proposition.

By a simple computation, one gets the following.

Lemma 4.3.3. *If (j,ℓ,t,u,v) is a permutation of $(1,2,3,4,5)$, then the following monomials are strictly inadmissible:*

i) $x_j^2x_\ell x_t x_u^3,\ j < \ell < t$.
ii) $x_j^2x_\ell x_t x_u x_v^2,\ j < \ell < t < u$.
iii) $x_1x_2^2x_3^2x_4x_5$.

The following is a corollary of a result in [24].

Lemma 4.3.4. *If x is one of the following monomials then $f_i(x)$, $1 \leqslant i \leqslant 5$, are strictly inadmissible:*

$$x_1^3x_2^{28}x_3x_4^3,\ x_1^3x_2^{28}x_3^3x_4,\ x_1^3x_2^7x_3^{24}x_4,\ x_1^7x_2^3x_3^{24}x_4,\ x_1^3x_2^5x_3^{25}x_4^2,\ x_1^3x_2^4x_3^{25}x_4^3,$$
$$x_1^3x_2^5x_3^{24}x_4^3,\ x_1^3x_2^4x_3^9x_4^{19},\ x_1^3x_2^4x_3^{11}x_4^{17},\ x_1^3x_2^5x_3^8x_4^{19},\ x_1^3x_2^5x_3^9x_4^{18},\ x_1^3x_2^5x_3^{10}x_4^{17},$$
$$x_1^3x_2^5x_3^{11}x_4^{16},\ x_1^3x_2^7x_3^8x_4^{17},\ x_1^7x_2^3x_3^8x_4^{17},\ x_1^3x_2^7x_3^9x_4^{16},\ x_1^7x_2^3x_3^9x_4^{16}.$$

Lemma 4.3.5. *The following monomials are strictly inadmissible:*

$$x_1x_2^6x_3^3x_4^8x_5^{17}\ \ x_1x_2^6x_3^3x_4^9x_5^{16}\ \ x_1x_2^6x_3^3x_4^{24}x_5\ \ x_1^3x_2^4x_3x_4^8x_5^{19}\ \ x_1^3x_2^4x_3x_4^9x_5^{18}$$
$$x_1^3x_2^4x_3x_4^{10}x_5^{17}\ \ x_1^3x_2^4x_3x_4^{11}x_5^{16}\ \ x_1^3x_2^4x_3x_4^{24}x_5^3\ \ x_1^3x_2^4x_3x_4^{25}x_5^2\ \ x_1^3x_2^4x_3^3x_4^8x_5^{17}$$
$$x_1^3x_2^4x_3^3x_4^9x_5^{16}\ \ x_1^3x_2^4x_3^3x_4^{24}x_5\ \ x_1^3x_2^4x_3^8x_4x_5^{19}\ \ x_1^3x_2^4x_3^8x_4^3x_5^{17}\ \ x_1^3x_2^4x_3^8x_4^{17}x_5^3$$
$$x_1^3x_2^4x_3^8x_4^{19}x_5\ \ x_1^3x_2^4x_3^9x_4x_5^{18}\ \ x_1^3x_2^4x_3^9x_4^2x_5^{17}\ \ x_1^3x_2^4x_3^9x_4^3x_5^{16}\ \ x_1^3x_2^4x_3^9x_4^{16}x_5^3$$
$$x_1^3x_2^4x_3^9x_4^{17}x_5^2\ \ x_1^3x_2^4x_3^9x_4^{18}x_5\ \ x_1^3x_2^4x_3^{11}x_4x_5^{16}\ \ x_1^3x_2^4x_3^{11}x_4^{16}x_5\ \ x_1^3x_2^4x_3^{24}x_4x_5^3$$
$$x_1^3x_2^4x_3^{24}x_4^3x_5\ \ x_1^3x_2^4x_3^{25}x_4x_5^2\ \ x_1^3x_2^4x_3^{25}x_4^2x_5\ \ x_1^3x_2^5x_3x_4^8x_5^{18}\ \ x_1^3x_2^5x_3x_4^{10}x_5^{16}$$
$$x_1^3x_2^5x_3x_4^{24}x_5^2\ \ x_1^3x_2^5x_3^8x_4x_5^{18}\ \ x_1^3x_2^5x_3^8x_4^2x_5^{17}\ \ x_1^3x_2^5x_3^8x_4^3x_5^{16}\ \ x_1^3x_2^5x_3^8x_4^{16}x_5^3$$
$$x_1^3x_2^5x_3^8x_4^{17}x_5^2\ \ x_1^3x_2^5x_3^8x_4^{18}x_5\ \ x_1^3x_2^5x_3^9x_4^2x_5^{16}\ \ x_1^3x_2^5x_3^9x_4^{16}x_5^2\ \ x_1^3x_2^5x_3^{10}x_4x_5^{16}$$
$$x_1^3x_2^5x_3^{10}x_4^{16}x_5\ \ x_1^3x_2^5x_3^{24}x_4x_5^2\ \ x_1^3x_2^5x_3^{24}x_4^2x_5\ \ x_1^3x_2^7x_3^8x_4x_5^{16}\ \ x_1^3x_2^7x_3^8x_4^{16}x_5$$
$$x_1^3x_2^{28}x_3x_4x_5^2\ \ x_1^3x_2^{28}x_3x_4^2x_5\ \ x_1^7x_2^3x_3^8x_4x_5^{16}\ \ x_1^7x_2^3x_3^8x_4^{16}x_5.$$

Proof. We prove the lemma for the monomials $x = x_1x_2^6x_3^3x_4^8x_5^{17}$, $y = x_1^3x_2^4x_3^3x_4^9x_5^{16}$. The others can be proved by the similar computations. By a direct computation, we have

$$\begin{aligned}
x &= x_1x_2^3x_3^5x_4^2x_5^{24} + x_1x_2^3x_3^5x_4^8x_5^{18} + x_1x_2^3x_3^6x_4x_5^{24} + x_1x_2^3x_3^6x_4^8x_5^{17} + x_1x_2^3x_3^8x_4x_5^{22}\\
&\quad + x_1x_2^3x_3^8x_4^2x_5^{21} + x_1x_2^4x_3^2x_4x_5^{27} + x_1x_2^4x_3^3x_4x_5^{26} + x_1x_2^4x_3^3x_4^2x_5^{25} + x_1x_2^4x_3^{10}x_4x_5^{19}\\
&\quad + x_1x_2^6x_3^2x_4x_5^{25} + x_1x_2^6x_3^3x_4x_5^{24} + Sq^1(x_1^2x_2^5x_3^5x_4x_5^{21}) + Sq^2(x_1x_2^6x_3^3x_4^2x_5^{21}\\
&\quad + x_1x_2^5x_3^5x_4x_5^{21} + x_1x_2^3x_3^6x_4x_5^{22} + x_1x_2^3x_3^6x_4^2x_5^{21} + x_1x_2^3x_3^5x_4^2x_5^{22} + x_1x_2^6x_3^3x_4x_5^{22}\\
&\quad + x_1x_2^6x_3^6x_4x_5^{19} + x_1x_2^6x_3^2x_4x_5^{23}) + Sq^4(x_1x_2^{10}x_3^3x_4^4x_5^{13} + x_1x_2^4x_3^3x_4^2x_5^{21}\\
&\quad + x_1x_2^3x_3^{10}x_4^4x_5^{13} + x_1x_2^3x_3^9x_4^4x_5^{14} + x_1x_2^4x_3^3x_4x_5^{22} + x_1x_2^{10}x_3^4x_4x_5^{15}\\
&\quad + x_1x_2^4x_3^2x_4x_5^{23} + x_1x_2^4x_3^6x_4x_5^{19}) + Sq^8(x_1x_2^6x_3^3x_4^4x_5^{13} + x_1x_2^3x_3^6x_4^4x_5^{13}\\
&\quad + x_1x_2^3x_3^5x_4^4x_5^{14} + x_1x_2^6x_3^4x_4x_5^{15}) \mod\big(P_5^-(\omega_{(1)})\big)
\end{aligned}$$

Hence, x is strictly inadmissible. By a similar computation, we obtain

$$\begin{aligned}
y &= x_1^2x_2x_3^2x_4^5x_5^{25} + x_1^2x_2x_3^3x_4^9x_5^{20} + x_1^2x_2x_3^3x_4^{12}x_5^{17} + x_1^2x_2x_3^4x_4^3x_5^{25}\\
&\quad + x_1^2x_2x_3^5x_4^9x_5^{18} + x_1^2x_2x_3^5x_4^{10}x_5^{17} + x_1^2x_2x_3^{10}x_4^5x_5^{17} + x_1^2x_2x_3^{12}x_4^3x_5^{17}\\
&\quad + x_1^3x_2x_3^2x_4^4x_5^{25} + x_1^3x_2x_3^2x_4^5x_5^{24} + x_1^3x_2x_3^4x_4^3x_5^{24} + x_1^3x_2x_3^4x_4^{10}x_5^{17}\\
&\quad + x_1^3x_2x_3^5x_4^{10}x_5^{16} + x_1^3x_2x_3^8x_4^3x_5^{20} + x_1^3x_2x_3^8x_4^5x_5^{18} + x_1^3x_2x_3^{10}x_4^4x_5^{17}\\
&\quad + x_1^3x_2^2x_3^4x_4^9x_5^{17} + x_1^3x_2^2x_3^5x_4^8x_5^{17} + x_1^3x_2^2x_3^8x_4^5x_5^{17} + x_1^3x_2^4x_3^2x_4^9x_5^{17}\\
&\quad + x_1^3x_2^4x_3^3x_4^8x_5^{17} + Sq^1(q_1) + Sq^2(q_2) + Sq^4(q_3) + Sq^8(q_4) \mod\big(P_5^-(\omega_{(1)})\big),
\end{aligned}$$

where

$$\begin{aligned}
q_1 &= x_1^3x_2x_3^2x_4^3x_5^{25} + x_1^3x_2x_3^3x_4^9x_5^{18} + x_1^3x_2x_3^3x_4^{10}x_5^{17}\\
&\quad + x_1^3x_2x_3^5x_4^5x_5^{20} + x_1^3x_2x_3^{10}x_4^3x_5^{17} + x_1^3x_2^4x_3^5x_4^5x_5^{17},\\
q_2 &= x_1^2x_2x_3^2x_4^3x_5^{25} + x_1^2x_2x_3^3x_4^9x_5^{18} + x_1^2x_2x_3^3x_4^{10}x_5^{17} + x_1^2x_2x_3^{10}x_4^3x_5^{17} + x_1^5x_2x_3^2x_4^3x_5^{22}\\
&\quad + x_1^5x_2x_3^6x_4^3x_5^{18} + x_1^5x_2^2x_3^2x_4^3x_5^{21} + x_1^5x_2^2x_3^3x_4^5x_5^{18} + x_1^5x_2^2x_3^3x_4^6x_5^{17} + x_1^5x_2^2x_3^6x_4^3x_5^{17},
\end{aligned}$$

$$\begin{aligned}q_3 &= x_1^3x_2x_3^2x_4^3x_5^{22} + x_1^3x_2x_3^6x_4^3x_5^{18} + x_1^3x_2x_3^8x_4^5x_5^{14} + x_1^3x_2x_3^9x_4^6x_5^{12}\\ &\quad + x_1^3x_2^2x_3^2x_4^3x_5^{21} + x_1^3x_2^2x_3^3x_4^5x_5^{18} + x_1^3x_2^2x_3^3x_4^6x_5^{17} + x_1^3x_2^2x_3^6x_4^3x_5^{17}\\ &\quad + x_1^3x_2^2x_3^8x_4^5x_5^{13} + x_1^3x_2^8x_3^2x_4^5x_5^{13} + x_1^3x_2^8x_3^3x_4^5x_5^{12} + x_1^3x_2^8x_3^4x_4^3x_5^{13}\\ q_4 &= x_1^3x_2x_3^4x_4^5x_5^{14} + x_1^3x_2x_3^5x_4^6x_5^{12} + x_1^3x_2^2x_3^4x_4^5x_5^{13}\\ &\quad + x_1^3x_2^4x_3^2x_4^5x_5^{13} + x_1^3x_2^4x_3^3x_4^5x_5^{12} + x_1^3x_2^4x_3^4x_4^3x_5^{13}.\end{aligned}$$

Hence, x is strictly inadmissible. □

Proof of Proposition 4.3.2. Let x be an admissible monomial such that $\omega(x) = \omega_{(1)}$. Then, $x = x_jx_\ell x_ty^2$ with $1 \leqslant j < \ell < t \leqslant 5$ and $y \in B_5(2,1,1,1)$.

Let $z \in B_5(2,1,1,1)$ such that $x_jx_\ell x_tz^2 \in P_5^+$. By a direct computation using the results in Subsection 4.2, we see that if $x_jx_\ell x_tz^2 \neq a_t, \forall t,\ 1 \leqslant t \leqslant 160$, then there is a monomial w which is given in one of Lemmas 4.3.3, 4.3.4 and 4.3.5 such that $x_jx_\ell x_tz^2 = wz_1^{2^u}$ with suitable monomial $z_1 \in P_5$, and $u = \max\{j \in \mathbb{Z} : \omega_j(w) > 0\}$. By Theorem 2.8, $x_jx_\ell x_tz^2$ is inadmissible. Since $x = x_jx_\ell x_ty^2$ and x is admissible, one gets $x = a_t$ for some $t,\ 1 \leqslant t \leqslant 160$. This implies $B_5^+(\omega_{(1)}) \subset \{a_t :\ 1 \leqslant t \leqslant 160\}$.

We now prove the set $\{[a_t] :\ 1 \leqslant t \leqslant 160\}$ is linearly independent in $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{35}$. Suppose there is a linear relation

$$\mathcal{S} = \sum_{t=1}^{160} \gamma_t a_t \equiv 0,$$

where $\gamma_t \in \mathbb{F}_2$. For $1 \leqslant i < j \leqslant 5$, we explicitly compute $p_{(i;j)}(\mathcal{S})$ in terms of the admissible monomials in P_4 $(\mathrm{mod}(\mathcal{A}^+P_4))$. By a direct computation from the relations $p_{(i;j)}(\mathcal{S}) \equiv 0$ with $1 \leqslant i < j \leqslant 5$, we obtain $\gamma_t = 0$ for $1 \leqslant t \leqslant 160$. The proposition follows. □

Proposition 4.3.6. $QP_5(\omega_{(2)}) = 0$.

We need the following lemma.

Lemma 4.3.7. *All permutations of the following monomials are strictly inadmissible:*

$$x_1^3x_2^4x_3^8x_4^9x_5^{11},\ x_1^3x_2^4x_3^9x_4^9x_5^{10},\ x_1^3x_2^5x_3^8x_4^8x_5^{11},\ x_1^3x_2^5x_3^8x_4^9x_5^{10},\ x_1^3x_2^7x_3^8x_4^8x_5^9.$$

Proof. We prove the lemma for the monomial $x = x_1^3x_2^4x_3^8x_4^9x_5^{11}$. The others can be proved by the similar computations. By a direct computation, we have

$$\begin{aligned}x &= Sq^1(x_1^3x_2x_3^2x_4^9x_5^{19}) + Sq^2(x_1^5x_2^2x_3^2x_4^5x_5^{19} + x_1^5x_2x_3^2x_4^6x_5^{19})\\ &\quad + Sq^4(x_1^3x_2^8x_3^4x_4^5x_5^{11} + x_1^3x_2^2x_3^2x_4^5x_5^{19} + x_1^3x_2^8x_3^2x_4^5x_5^{13} + x_1^3x_2^2x_3^8x_4^5x_5^{13}\\ &\quad + x_1^3x_2x_3^2x_4^6x_5^{19} + x_1^3x_2x_3^8x_4^6x_5^{13}) + Sq^8(x_1^3x_2^4x_3^4x_4^5x_5^{11} + x_1^3x_2^4x_3^2x_4^5x_5^{13}\\ &\quad + x_1^3x_2^2x_3^4x_4^5x_5^{13} + x_1^3x_2x_3^4x_4^6x_5^{13}) \mod\big(P_5^-(\omega_{(2)})\big)\end{aligned}$$

This equality shows that all permutations of x are strictly inadmissible. □

Proof of Proposition 4.3.6. Let x be an admissible monomial such that $\omega(x) = \omega_{(2)}$. Then $x = x_jx_\ell x_ty^2$ with $y \in B_5(2,1,3)$.

Let $z \in B_5(2,1,3)$ such that $x_jx_\ell x_tz^2 \in P_5^+$. By a direct computation using the results in Subsection 4.2, we see that if $x_jx_\ell x_tz^2$ is not a permutation of one of monomials as given in Lemma 4.3.7, then there is a monomial w which is given

in Lemma 4.3.3 such that $x_jx_\ell x_tz^2 = wz_1^{2^u}$ with suitable monomial $z_1 \in P_5$, and $u = \max\{j \in \mathbb{Z} : \omega_j(w) > 0\}$. By Theorem 2.8, $x_jx_\ell x_tz^2$ is inadmissible. Since $x = x_jx_\ell x_ty^2$ and x is admissible, x is a permutation of one of monomials as given in Lemma 4.3.7. Now the proposition follows from Lemma 4.3.7. □

Proposition 4.3.8. $QP_5(\omega_{(3)}) = 0$.

The following lemma is needed for the proof of the proposition.

Lemma 4.3.9. *The following monomials are strictly inadmissible:*
i) $x_j^3x_\ell^4x_t^5x_u^7,\ x_j^3x_\ell^5x_t^5x_u^6$.
ii) $x_j^3x_\ell^4x_tx_u^4x_v^7,\ x_j^3x_\ell^4x_tx_u^5x_v^6,\ x_j^3x_\ell^4x_t^2x_u^5x_v^5,\ j < \ell < t$; $x_j^3x_\ell^4x_t^3x_u^4x_v^5,\ j < \ell,\ t > 3$.
Here (j, ℓ, t, u, v) *is a permutation of* $(1, 2, 3, 4, 5)$.
iii) *All permutations of the monomials:*

$$\begin{array}{llll}
x_1x_2^2x_3^7x_4^{12}x_5^{13} & x_1x_2^3x_3^6x_4^{12}x_5^{13} & x_1x_2^3x_3^7x_4^{12}x_5^{12} & x_1x_2^6x_3^{11}x_4^4x_5^{13}\\
x_1x_2^7x_3^{10}x_4^4x_5^{13} & x_1x_2^7x_3^{11}x_4^4x_5^{12} & x_1x_2^6x_3^{11}x_4^5x_5^{12} & x_1x_2^7x_3^{10}x_4^5x_5^{12}\\
x_1^3x_2^5x_3^2x_4^{12}x_5^{13} & x_1^3x_2^3x_3^4x_4^{12}x_5^{13} & x_1^3x_2^3x_3^5x_4^{12}x_5^{12} & x_1^3x_2^4x_3^{11}x_4^4x_5^{13}\\
x_1^3x_2^5x_3^{10}x_4^4x_5^{13} & x_1^3x_2^4x_3^{11}x_4^5x_5^{12} & x_1^3x_2^7x_3^9x_4^4x_5^{12} & x_1^3x_2^5x_3^{10}x_4^5x_5^{12}.
\end{array}$$

Proof. We prove the lemma for $x = x_1x_2^2x_3^7x_4^{12}x_5^{13}$ and $y = x_1x_2^3x_3^6x_4^{12}x_5^{13}$. The others are proved by the similar computations. A direct computation shows

$$\begin{aligned}
x &= Sq^1(x_1^2x_2x_3^7x_4^5x_5^{19} + x_1^2x_2x_3^9x_4^3x_5^{19} + x_1^2x_2x_3^7x_4^3x_5^{21})\\
&\quad + Sq^2(x_1x_2^4x_3^7x_4^{10}x_5^{11} + x_1x_2x_3^7x_4^5x_5^{19} + x_1x_2x_3^9x_4^3x_5^{19} + x_1x_2x_3^7x_4^3x_5^{21})\\
&\quad + Sq^4(x_1x_2^2x_3^{11}x_4^6x_5^{11} + x_1x_2^2x_3^6x_4^3x_5^{19} + x_1x_2^2x_3^{11}x_4^3x_5^{14})\\
&\quad + Sq^8(x_1x_2^2x_3^7x_4^6x_5^{11} + x_1x_2^2x_3^7x_4^3x_5^{14}) \mod\big(P_5^-(\omega_{(3)})\big)
\end{aligned}$$

Hence, all permutations of x are strictly inadmissible. We have

$$\begin{aligned}
y &= Sq^1\big(x_1x_2^3x_3^2x_4^9x_5^{19} + x_1x_2^3x_3^3x_4^{12}x_5^{15} + x_1x_2^3x_3^5x_4^{10}x_5^{15}\\
&\quad + x_1x_2^3x_3^9x_4^2x_5^{19} + x_1x_2^3x_3^{10}x_4^5x_5^{15} + x_1x_2^3x_3^{12}x_4^3x_5^{15} + x_1x_2^6x_3^3x_4^3x_5^{21}\\
&\quad + x_1x_2^8x_3^3x_4^3x_5^{19} + x_1x_2^{12}x_3^3x_4^3x_5^{15} + x_1^2x_2^5x_3^3x_4^9x_5^{15} + x_1^2x_2^5x_3^9x_4^3x_5^{15}\\
&\quad + x_1^4x_2^3x_3^3x_4^9x_5^{15} + x_1^4x_2^3x_3^9x_4^3x_5^{15} + x_1^4x_2^5x_3^3x_4^3x_5^{19} + x_1^4x_2^9x_3^3x_4^3x_5^{15}\big)\\
&\quad + Sq^2\big(x_1x_2^2x_3^3x_4^{12}x_5^{15} + x_1x_2^2x_3^5x_4^{10}x_5^{15} + x_1x_2^4x_3^{10}x_4^3x_5^{15}\\
&\quad + x_1x_2^5x_3^2x_4^{10}x_5^{15} + x_1x_2^5x_3^6x_4^6x_5^{15} + x_1x_2^5x_3^6x_4^{10}x_5^{11} + x_1x_2^5x_3^{10}x_4^2x_5^{15}\\
&\quad + x_1x_2^{10}x_3^2x_4^5x_5^{15} + x_1x_2^{10}x_3^5x_4^2x_5^{15} + x_1^2x_2^2x_3^5x_4^9x_5^{15} + x_1^2x_2^2x_3^9x_4^5x_5^{15}\\
&\quad + x_1^2x_2^3x_3^3x_4^{10}x_5^{15} + x_1^2x_2^3x_3^{10}x_4^3x_5^{15} + x_1^2x_2^5x_3^2x_4^9x_5^{15} + x_1^2x_2^5x_3^9x_4^2x_5^{15}\\
&\quad + x_1^2x_2^6x_3^3x_4^3x_5^{19} + x_1^2x_2^9x_3^2x_4^5x_5^{15} + x_1^2x_2^9x_3^5x_4^2x_5^{15} + x_1^2x_2^{10}x_3^3x_4^3x_5^{15}\\
&\quad + x_1^4x_2^3x_3^2x_4^9x_5^{15} + x_1^4x_2^3x_3^9x_4^2x_5^{15} + x_1^4x_2^6x_3^3x_4^5x_5^{15} + x_1^4x_2^6x_3^5x_4^3x_5^{15}\big)\\
&\quad + Sq^4\big(x_1x_2^3x_3^2x_4^{10}x_5^{15} + x_1x_2^3x_3^6x_4^6x_5^{15} + x_1x_2^3x_3^{10}x_4^2x_5^{15}\\
&\quad + x_1x_2^3x_3^{10}x_4^6x_5^{11} + x_1x_2^6x_3^2x_4^3x_5^{19} + x_1x_2^6x_3^3x_4^2x_5^{19} + x_1x_2^{10}x_3^2x_4^3x_5^{15}\\
&\quad + x_1x_2^{10}x_3^3x_4^2x_5^{15} + x_1x_2^{10}x_3^3x_4^3x_5^{14} + x_1^2x_2^2x_3^3x_4^5x_5^{19} + x_1^2x_2^2x_3^3x_4^9x_5^{15}
\end{aligned}$$

$$+ x_1^2x_2^2x_3^5x_4^3x_5^{19} + x_1^2x_2^2x_3^9x_4^3x_5^{15} + x_1^2x_2^3x_3^2x_4^5x_5^{19} + x_1^2x_2^3x_3^3x_4^6x_5^{17}$$
$$+ x_1^2x_2^3x_3^5x_4^2x_5^{19} + x_1^2x_2^3x_3^6x_4^3x_5^{17} + x_1^2x_2^5x_3^2x_4^3x_5^{19} + x_1^2x_2^5x_3^3x_4^2x_5^{19}$$
$$+ x_1^2x_2^6x_3^3x_4^3x_5^{17} + x_1^2x_2^6x_3^3x_4^5x_5^{15} + x_1^2x_2^6x_3^5x_4^3x_5^{15} + x_1^2x_2^9x_3^2x_4^3x_5^{15}$$
$$+ x_1^2x_2^9x_3^3x_4^2x_5^{15}\big) + Sq^8\big(x_1x_2^3x_3^6x_4^6x_5^{11} + x_1x_2^6x_3^3x_4^3x_5^{14}\big) \mod\big(P_5^-(\omega_{(3)})\big).$$

This equality implies that all permutations of y are strictly inadmissible. □

Proof of Proposition 4.3.8. Let x be an admissible monomial such that $\omega(x) = \omega_{(3)}$. Then $x = x_jx_\ell x_t y^2$ with $1 \leqslant j < \ell < t \leqslant 5$ and $y \in B_5(2,3,2)$. By a direct computation using Proposition 4.2.2, we see that there is a monomial w which is given in one of Lemmas 4.3.3, 4.3.9 such that $x_jx_\ell x_t y^2 = wz_1^{2^u}$ with suitable monomial $z_1 \in P_5$, and $u = \max\{j \in \mathbb{Z} : \omega_j(w) > 0\}$. By Theorem 2.8, $x = x_jx_\ell x_t y^2$ is inadmissible. Hence, $QP_5(\omega_{(3)}) = 0$. □

Consider the monomials $a_t = a_{35,t},\ 161 \leqslant t \leqslant 210$ as given in Subsection 6.5.

Proposition 4.3.10. *The $\mathbb{F}_2$-vector space $QP_5(\omega_{(4)})$ is an GL_5-module generated by the class $[a_{203}]_{\omega_{(4)}}$ and $B_5(\omega_{(4)}) = \{a_t : 161 \leqslant t \leqslant 210\}$. Consequently,* $\dim QP_5(\omega_{(4)}) = 50$.

We need the following lemmas.

Lemma 4.3.11. *The following monomials are strictly inadmissible:*
i) $x_j^2x_\ell x_t^2x_u^3x_v^3,\ j < \ell;\ x_j^2x_\ell^3x_t^3x_u^3$.
Here (j,ℓ,t,u,v) is a permutation of $(1,2,3,4,5)$.
ii) *All permutations of the monomials:*

$$\begin{array}{llll} x_1x_2^2x_3^2x_4^{15}x_5^{15} & x_1x_2^2x_3^3x_4^{14}x_5^{15} & x_1x_2^2x_3^7x_4^{10}x_5^{15} & x_1x_2^3x_3^3x_4^{14}x_5^{14} \\ x_1x_2^3x_3^6x_4^{10}x_5^{15} & x_1x_2^7x_3^7x_4^{10}x_5^{10} & x_1^2x_2^3x_3^3x_4^{13}x_5^{14} & x_1^2x_2^3x_3^5x_4^{10}x_5^{15} \\ x_1^2x_2^2x_3^3x_4^{13}x_5^{15} & x_1^2x_2^2x_3^7x_4^9x_5^{15} & x_1^2x_2^7x_3^7x_4^9x_5^{10}. & \end{array}$$

Proof. We prove the lemma for $x = x_1x_2^2x_3^7x_4^{10}x_5^{15}$, $y = x_1x_2^3x_3^3x_4^{14}x_5^{14}$ and $z = x_1x_2^7x_3^7x_4^{10}x_5^{10}$. We have

$$x = Sq^1(x_1^2x_2x_3^7x_4^9x_5^{15}) + Sq^2(x_1x_2x_3^7x_4^9x_5^{15} + x_1x_2x_3^3x_4^9x_5^{19} + x_1x_2x_3^7x_4^3x_5^{21}$$
$$+ x_1x_2x_3^3x_4^3x_5^{25}) + Sq^4(x_1x_2x_3^5x_4^9x_5^{15} + x_1x_2x_3^{11}x_4^5x_5^{13} + x_1x_2x_3^5x_4^3x_5^{21}$$
$$+ x_1x_2x_3^3x_4^5x_5^{21}) + Sq^8(x_1x_2x_3^7x_4^5x_5^{13}) \mod\big(P_5^-(\omega_{(4)})\big).$$

So, all permutations of x are strictly inadmissible.

$$y = Sq^1\big(x_1x_2^3x_3^{10}x_4^{13}x_5^7 + x_1^2x_2^3x_3^9x_4^{13}x_5^7 + x_1x_2^3x_3^3x_4^{14}x_5^{13} + x_1x_2^3x_3^5x_4^{14}x_5^{11}$$
$$+ x_1x_2^3x_3^3x_4^{13}x_5^{14}\big) + Sq^2\big(x_1x_2^5x_3^3x_4^7x_5^{17} + x_1x_2^5x_3^3x_4^{11}x_5^{13}$$
$$+ x_1x_2^5x_3^5x_4^{11}x_5^{11} + x_1x_2^5x_3^9x_4^{11}x_5^7\big) + Sq^4\big(x_1x_2^3x_3^3x_4^7x_5^{17}$$
$$+ x_1x_2^3x_3^3x_4^{11}x_5^{13} + x_1x_2^3x_3^5x_4^{11}x_5^{11} + x_1x_2^3x_3^9x_4^{11}x_5^7 + x_1x_2^3x_3^5x_4^{13}x_5^9$$
$$+ x_1^2x_2^3x_3^6x_4^{13}x_5^7\big) + Sq^8\big(x_1x_2^3x_3^5x_4^9x_5^9\big),\ \mod\big(P_5^-(\omega_{(4)})\big).$$

Hence, all permutations of y are strictly inadmissible.

$$\begin{aligned}z = {} & Sq^1\big(x_1x_2^7x_3^7x_4^9x_5^{10} + x_1x_2^7x_3^{11}x_4^3x_5^{12} + x_1x_2^7x_3^{13}x_4^3x_5^{10} + x_1x_2^{11}x_3^7x_4^3x_5^{12}\\ & + x_1x_2^{13}x_3^7x_4^3x_5^{10} + x_1^2x_2^7x_3^{13}x_4^3x_5^9 + x_1^2x_2^{13}x_3^7x_4^3x_5^9 + x_1^4x_2^7x_3^{11}x_4^3x_5^9\\ & + x_1^4x_2^{11}x_3^7x_4^3x_5^9\big) + Sq^2\big(x_1x_2^7x_3^{11}x_4^5x_5^9 + x_1x_2^{11}x_3^7x_4^5x_5^9 + x_1^2x_2^7x_3^{11}x_4^3x_5^{10}\\ & + x_1^2x_2^{11}x_3^7x_4^3x_5^{10}\big) + Sq^4\big(x_1x_2^7x_3^{11}x_4^3x_5^9 + x_1x_2^9x_3^{13}x_4^3x_5^5 + x_1x_2^{11}x_3^7x_4^3x_5^9\\ & + x_1x_2^{13}x_3^9x_4^3x_5^5 + x_1^2x_2^7x_3^7x_4^5x_5^{10}\big) + Sq^8\big(x_1x_2^9x_3^9x_4^3x_5^5\big), \mod\big(P_5^-(\omega_{(4)})\big).\end{aligned}$$

Hence, all permutations of z are strictly inadmissible. □

Lemma 4.3.12. *The following monomials are strictly inadmissible:*

$$\begin{array}{llllll}
x_1x_2^3x_3^6x_4^{14}x_5^{11} & x_1x_2^3x_3^{14}x_4^6x_5^{11} & x_1x_2^3x_3^{14}x_4^7x_5^{10} & x_1x_2^7x_3^{10}x_4^3x_5^{14} & x_1^3x_2x_3^6x_4^{14}x_5^{11}\\
x_1^3x_2x_3^{14}x_4^6x_5^{11} & x_1^3x_2x_3^{14}x_4^7x_5^{10} & x_1^3x_2^5x_3^2x_4^{14}x_5^{11} & x_1^3x_2^5x_3^6x_4^{10}x_5^{11} & x_1^3x_2^5x_3^6x_4^{11}x_5^{10}\\
x_1^3x_2^5x_3^7x_4^{10}x_5^{10} & x_1^3x_2^5x_3^{10}x_4^3x_5^{14} & x_1^3x_2^5x_3^{10}x_4^6x_5^{11} & x_1^3x_2^5x_3^{10}x_4^7x_5^{10} & x_1^3x_2^5x_3^{10}x_4^{14}x_5^3\\
x_1^3x_2^5x_3^{14}x_4^2x_5^{11} & x_1^3x_2^5x_3^{14}x_4^3x_5^{10} & x_1^3x_2^5x_3^{14}x_4^{10}x_5^3 & x_1^3x_2^5x_3^{14}x_4^{11}x_5^2 & x_1^3x_2^{13}x_3^2x_4^6x_5^{11}\\
x_1^3x_2^{13}x_3^2x_4^7x_5^{10} & x_1^3x_2^{13}x_3^3x_4^6x_5^{10} & x_1^3x_2^{13}x_3^6x_4^2x_5^{11} & x_1^3x_2^{13}x_3^6x_4^3x_5^{10} & x_1^3x_2^{13}x_3^6x_4^{10}x_5^3\\
x_1^3x_2^{13}x_3^6x_4^{11}x_5^2 & x_1^3x_2^{13}x_3^7x_4^2x_5^{10} & x_1^3x_2^{13}x_3^7x_4^{10}x_5^2 & x_1^7x_2x_3^{10}x_4^3x_5^{14} & x_1^7x_2^9x_3^2x_4^3x_5^{14}\\
x_1^7x_2^9x_3^3x_4^2x_5^{14} & x_1^7x_2^9x_3^3x_4^6x_5^{10} & x_1^7x_2^9x_3^3x_4^{14}x_5^2. & &
\end{array}$$

Proof. We prove the lemma for $x = x_1x_2^3x_3^{14}x_4^7x_5^{10}$ and $y = x_1x_2^7x_3^{10}x_4^3x_5^{14}$. The others can be obtained by the similar computations. We have

$$\begin{aligned}x = {} & x_1x_2^3x_3^7x_4^{14}x_5^{10} + Sq^1\big(x_1^2x_2^3x_3^{13}x_4^7x_5^9 + x_1^2x_2^3x_3^7x_4^{13}x_5^9\big) + Sq^2\big(x_1x_2^5x_3^{11}x_4^7x_5^9\\ & + x_1x_2^5x_3^7x_4^{11}x_5^9\big) + Sq^4\big(x_1x_2^3x_3^{11}x_4^7x_5^9 + x_1x_2^3x_3^7x_4^{11}x_5^9 + x_1x_2^3x_3^{13}x_4^9x_5^5\\ & + x_1x_2^3x_3^9x_4^{13}x_5^5\big) + Sq^8\big(x_1x_2^3x_3^9x_4^9x_5^5\big), \mod\big(P_5^-(\omega_{(4)})\big).\end{aligned}$$

Hence, x is strictly inadmissible.

$$\begin{aligned}y = {} & x_1x_2^7x_3^3x_4^{10}x_5^{14} + Sq^1\big(x_1x_2^7x_3^9x_4^{10}x_5^7 + x_1x_2^7x_3^{10}x_4^3x_5^{13} + x_1x_2^7x_3^{10}x_4^9x_5^7\\ & + x_1x_2^7x_3^{12}x_4^3x_5^{11} + x_1x_2^{11}x_3^3x_4^{12}x_5^7 + x_1x_2^{13}x_3^3x_4^{10}x_5^7 + x_1^2x_2^7x_3^3x_4^9x_5^{13}\\ & + x_1^2x_2^7x_3^5x_4^9x_5^{11} + x_1^2x_2^{13}x_3^3x_4^9x_5^7 + x_1^4x_2^7x_3^9x_4^3x_5^{11} + x_1^4x_2^{11}x_3^3x_4^9x_5^7\big)\\ & + Sq^2\big(x_1x_2^7x_3^5x_4^9x_5^{11} + x_1x_2^{11}x_3^5x_4^9x_5^7 + x_1^2x_2^7x_3^{10}x_4^3x_5^{11} + x_1^2x_2^{11}x_3^3x_4^{10}x_5^7\big)\\ & + Sq^4\big(x_1x_2^7x_3^3x_4^9x_5^{11} + x_1x_2^9x_3^3x_4^5x_5^{13} + x_1x_2^{11}x_3^3x_4^9x_5^7\\ & + x_1x_2^{13}x_3^3x_4^5x_5^9 + x_1^2x_2^7x_3^5x_4^{10}x_5^7 + x_1^2x_2^{11}x_3^6x_4^5x_5^7\big)\\ & + Sq^8\big(x_1x_2^9x_3^3x_4^5x_5^9 + x_1^2x_2^7x_3^6x_4^5x_5^7\big), \mod\big(P_5^-(\omega_{(4)})\big).\end{aligned}$$

This equality implies that y are strictly inadmissible. □

Lemma 4.3.13. *The class $[a_{161}]_{\omega_{(4)}}$ is non-zero in the vector space $QP_5(\omega_{(4)})$.*

Proof. Suppose the contrary that $[a_{161}]_{\omega_{(4)}} = 0$. Then

$$a_{161} = \sum_{u=0}^{4} Sq^{2^u}(B_u) \mod(P_5^-(\omega_{(4)})), \tag{4.1}$$

where B_u are suitable polynomials in P_5. We set

$$f_{4,5} = x_1^2x_2^4x_3^7x_4^{14}x_5^{14} + x_1^4x_2^2x_3^7x_4^{14}x_5^{14},\ y_1 = x_1x_2x_3^7x_4^7x_5^{11},$$
$$f_{3,5} = x_1^2x_2^4x_3^{14}x_4^7x_5^{14} + x_1^4x_2^2x_3^{14}x_4^7x_5^{14},\ y_2 = x_1x_2x_3^7x_4^{11}x_5^7,$$
$$f_{3,4} = x_1^2x_2^4x_3^{14}x_4^{14}x_5^7 + x_1^4x_2^2x_3^{14}x_4^{14}x_5^7,\ y_3 = x_1x_2x_3^{11}x_4^7x_5^7.$$

By a direct computation, we can see that there are polynomial $\bar{B}_u,\ u = 0, 1, 2$ such that

$$Sq^8(y_t) = \sum_{u=0}^{2} Sq^{2^u}(\bar{B}_u) \bmod(P_5^-(\omega_{(4)}) + \mathcal{B})) \tag{4.2}$$

where $\mathcal{B}$ is the subspace of $(P_5)_{35}$ spanned by all monomials x such that $\max\{\nu_i(x) : 1 \leqslant i \leqslant 5\} > 14$. By combining (4.1) and (4.1), one gets

$$a_{161} = \sum_{u=0}^{4} Sq^{2^u}(A_u) + \ \bmod(P_5^-(\omega_{(4)}) + \mathcal{B}), \tag{4.3}$$

where A_u are suitable polynomials such that y_t is not a term of A_3 with $t = 1, 2, 3$ (recall that a monomial x in P_k is called *a term* of a polynomial f if it appears in the expression of f in terms of the monomial basis of P_k) .

Let $(Sq^2)^3$ acts on the both sides of (4.3). Observe that if x is a monomial in $P_5^-(\omega_{(4)})$ then, $(Sq^2)^3(x) \in P_5^-(1, 4, 4, 2)$ and $(Sq^2)^3Sq^1 = (Sq^2)^3Sq^2 = 0$. So, we get

$$(Sq^2)^3(a_{161}) = \sum_{u=2}^{4}(Sq^2)^3(Sq^{2^u}(A_u)) \bmod(P_5^-(1, 4, 4, 2) + (Sq^2)^3(\mathcal{B})).$$

It is easy to see that $f_{i,j} \notin P_5^-(1, 4, 4, 2)+(Sq^2)^3(\mathcal{B})$ and $f_{4,5}$ is a term of $(Sq^2)^3(a_{161})$ (recall that a polynomial g in P_k is called *a term* of a polynomial f if a monomial x is a term of g, then it is also a term of f).

By a direct computation, we see that $f_{i,j}$ is not a term of $(Sq^2)^3(Sq^{16}(A_4))$ for any $A_4 \in (P_5)_{19}$. If $f_{i,j}$ is a term of $(Sq^2)^3(Sq^8(A_3))$, then y_t is a term of A_3 and $f_{i,j}$ is a term of $(Sq^2)^3(Sq^8(y_t))$ with some $t,\ t = 1, 2, 3$. However, the polynomial $f_{i,j}$ is not a term of $(Sq^2)^3(Sq^8(y_t))$.

Since $f_{4,5}$ is a term of $(Sq^2)^3(a_{161})$, it must be a term of $(Sq^2)^3(Sq^4(A_2))$. We observe that $f_{4,5}$ is a term of $(Sq^2)^3(p_{1,j}),\ 1 \leqslant j \leqslant 8$, where

$$p_{1,1} = x_1x_2x_3^7x_4^{13}x_5^{13},\ p_{1,2} = x_1x_2^2x_3^7x_4^{11}x_5^{14},\ p_{1,3} = x_1x_2^2x_3^7x_4^{14}x_5^{11},$$
$$p_{1,4} = x_1^2x_2^2x_3^7x_4^{11}x_5^{13},\ p_{1,5} = x_1^2x_2^2x_3^7x_4^{13}x_5^{11},\ p_{1,6} = x_1x_2^3x_3^7x_4^{11}x_5^{13} + x_1^3x_2x_3^7x_4^{11}x_5^{13},$$
$$p_{1,7} = x_1x_2^3x_3^7x_4^{13}x_5^{11} + x_1^3x_2x_3^7x_4^{13}x_5^{11},\ p_{1,8} = x_1x_2^3x_3^7x_4^{11}x_5^{13} + x_1^3x_2x_3^7x_4^{13}x_5^{11}.$$

Hence, one of the following polynomials is a term of A_2:

$$q_{1,1} = x_1x_2x_3^7x_4^{11}x_5^{11},\ q_{1,2} = x_1x_2^2x_3^7x_4^7x_5^{14},\ q_{1,3} = x_1x_2^2x_3^7x_4^{14}x_5^7,$$
$$q_{1,4} = x_1^2x_2^2x_3^7x_4^7x_5^{13},\ q_{1,5} = x_1^2x_2^2x_3^7x_4^{13}x_5^7,\ q_{1,6} = x_1x_2^3x_3^7x_4^7x_5^{13} + x_1^3x_2x_3^7x_4^7x_5^{13},$$
$$q_{1,7} = x_1x_2^3x_3^7x_4^{13}x_5^7 + x_1^3x_2x_3^7x_4^{13}x_5^7,\ q_{1,8} = x_1x_2^3x_3^7x_4^{13}x_5^7 + x_1^3x_2x_3^7x_4^7x_5^{13},$$
$$q_{1,9} = x_1x_2^3x_3^7x_4^{13}x_5^7 + x_1^3x_2x_3^7x_4^7x_5^{13}.$$

Suppose $q_{1,9}$ is a term of A_2. Then $\bar{p} = x_1^2x_2^4x_3^{14}x_4^{14}x_5^7 + x_1^4x_2^2x_3^{14}x_4^7x_5^{14}$ is a term of $(Sq^2)^3(a_{161} + Sq^4(q_{1,9}))$. This implies $q_{2,9} = x_1^3x_2x_3^{13}x_4^7x_5^7 + x_1x_2^3x_3^{13}x_4^7x_5^7$ is a term of A_2. Then, $\tilde{p} = x_1^4x_2^2x_3^{14}x_4^{14}x_5^7 + x_1^2x_2^4x_3^{14}x_4^7x_5^{14}$ is a term of $(Sq^2)^3(a_{161} +$

$Sq^4(q_{1,9}+q_{2,9}))$. Hence, $q_{3,9}=x_1x_2^3x_3^7x_4^{13}x_5^7+x_1^3x_2x_3^7x_4^7x_5^{13}$ is a term of A_2. Now, $f_{4,5}$ is a term of $(Sq^2)^3(a_{161}+Sq^4(q_{1,9}+q_{2,9}+q_{3,9}))$ and $q_{i,9}, i=1,2,3$, are not the terms of $A_2+q_{1,9}+q_{2,9}+q_{3,9}$.

Now, we assume that $q_{i,9}, i=1,2,3$, and $q_{1,1}$ are the terms of A_2. Then, $y^*=x_1x_2x_3^{11}x_4^{11}x_5^{11}$ is a term of $a_{161}+Sq^4(q_{1,1})$, hence y^* is a term of $Sq^4(A_2+q_{1,1})+Sq^8(A_3)+Sq^{16}(A_4)$. Since y_1,y_2,y_3 are not the terms of A_3, y^* is a term of $Sq^4(A_2)$. So, either $q_{2,1}=x_1x_2x_3^{11}x_4^7x_5^{11}$ or $q_{3,1}=x_1x_2x_3^{11}x_4^{11}x_5^7$ is a term of A_2. If $q_{2,1}$ is a term of A_2 then $f_{3,5}$ is a term of $(Sq^2)^3(a_{161}+Sq^4(q_{1,1}+q_{2,1}))$. By an argument analogous to the previous one, we see that one of the following polynomials is a term of A_2:

$$q_{2,1}=x_1x_2x_3^{11}x_4^7x_5^{11},\ q_{2,2}=x_1x_2^2x_3^7x_4^7x_5^{14},\ q_{2,3}=x_1x_2^2x_3^{14}x_4^7x_5^7,$$
$$q_{2,4}=x_1^2x_2^2x_3^7x_4^7x_5^{13},\ q_{2,5}=x_1^2x_2^2x_3^{13}x_4^7x_5^7,\ q_{2,6}=x_1x_2^3x_3^7x_4^7x_5^{13}+x_1^3x_2x_3^7x_4^7x_5^{13},$$
$$q_{2,7}=x_1x_2^3x_3^{13}x_4^7x_5^7+x_1^3x_2x_3^{13}x_4^7x_5^7,\ q_{2,8}=x_1x_2^3x_3^{13}x_4^7x_5^7+x_1^3x_2x_3^7x_4^7x_5^{13}.$$

If $q_{3,1}$ is a term of A_2, then $f_{3,4}$ is a term of $(Sq^2)^3(a_{161}+Sq^4(q_{1,1}+q_{3,1}))$. Hence, one of the following polynomials is a term of A_2:

$$q_{3,1}=x_1x_2x_3^{11}x_4^{11}x_5^7,\ q_{3,2}=x_1x_2^2x_3^7x_4^{14}x_5^7,\ q_{3,3}=x_1x_2^2x_3^{14}x_4^7x_5^7,$$
$$q_{3,4}=x_1^2x_2^2x_3^7x_4^{13}x_5^7,\ q_{3,5}=x_1^2x_2^2x_3^{13}x_4^7x_5^7,\ q_{3,6}=x_1x_2^3x_3^7x_4^{13}x_5^7+x_1^3x_2x_3^7x_4^{13}x_5^7,$$
$$q_{3,7}=x_1x_2^3x_3^{13}x_4^7x_5^7+x_1^3x_2x_3^{13}x_4^7x_5^7,\ q_{3,8}=x_1x_2^3x_3^{13}x_4^7x_5^7+x_1^3x_2x_3^7x_4^{13}x_5^7.$$

By repeating the above argument and after a finite number of steps, we see that $q_{i,j}$ is a term of A_2 for $i=1,2,3,\ 1\leqslant j\leqslant 9$. Then, $f_{4,5}$ is a term of $(Sq^2)^3(a_{161}+\sum_{(i,j)}q_{i,j})$. However, it is not a term of

$$(Sq^2)^3(Sq^4(A_2+\sum_{(i,j)}q_{i,j}))+(Sq^2)^3(Sq^8(A_3))+(Sq^2)^3(Sq^{16}(A_4)).$$

This is a contradiction. The lemma is proved. □

Proof of Proposition 4.3.10. Let x be an admissible monomial in $P_5(\omega_{(4)})$. Then $x=x_jx_\ell x_ty^2$ with $y\in B_5(4,2,2)$.

Let $z\in B_5(4,2,2)$. By a direct computation using the results in Subsection 4.2, we see that if $x_jx_\ell x_tz^2\neq a_t,\ 161\leqslant t\leqslant 210$, then there is a monomial w which is given in one of Lemmas 4.3.11 and 4.3.12 such that $x_jx_\ell x_tz^2=wz_1^{2^u}$ with suitable monomial $z_1\in P_5$, and $u=\max\{j\in\mathbb{Z}:\omega_j(w)>0\}$. By Theorem 2.8, $x_jx_\ell x_tz^2$ is inadmissible. Since $x=x_jx_\ell x_ty^2$, x is admissible and $y\in B_5(4,2,2)$, one gets $x=a_t$ for some $t,\ 161\leqslant t\leqslant 210$. This implies $B_5(\omega_{(4)})\subset\{a_t:\ 161\leqslant t\leqslant 210\}$.

By a direct computation, we see that there is a direct summand decomposition of the Σ_5-modules:

$$QP_5(\omega_{(4)})=\langle[\Sigma_5(a_{162})]_{\omega_{(4)}}\rangle\bigoplus\langle[\Sigma_5(a_{175})]_{\omega_{(4)}}\rangle\bigoplus\langle[\Sigma_5(a_{203})]_{\omega_{(4)}}\rangle.$$

Consider the homomorphis $g_i:P_5\to P_5,\ 1\leqslant i\leqslant 5$, as defined in Section 2. Let $g=g_2g_1g_2g_5g_2g_3g_1g_2$. Since $a_{162}\equiv_{\omega_{(4)}}g(a_{175})+a_{175}$ and $a_{175}\equiv_{\omega_{(4)}}g_5(a_{203})+a_{203}$, $QP_5(\omega_{(4)})$ is the GL_5-module generated by the class $[a_{203}]_{\omega_{(4)}}$.

We now prove the set $\{[a_t]_{\omega_{(4)}}:\ 161\leqslant t\leqslant 210\}$ is linearly independent in $QP_5(\omega_{(4)})$. Suppose there is a linear relation

$$\mathcal{S}=\sum_{t=161}^{210}\gamma_ta_t\equiv_{\omega_{(4)}}0,\tag{4.4}$$

where $\gamma_t \in \mathbb{F}_2$. We prove that $\gamma_t = 0,\ \forall t,\ 161 \leqslant t \leqslant 210$.

Set $\mathcal{S}_1 = g_5(\mathcal{S}) + \mathcal{S} \equiv_{\omega_{(4)}} 0$, $\mathcal{S}_2 = (g_5g_2g_3)(\mathcal{S}_1) + \mathcal{S}_1 \equiv_{\omega_{(4)}} 0$. A direct computation using (4.4) shows that

$$\mathcal{S}_3 := (g_5)(\mathcal{S}_2) + \mathcal{S}_2 \equiv_{\omega_{(4)}} \gamma_{181}a_{175} + \gamma_{183}a_{163} \equiv_{\omega_{(4)}} 0. \tag{4.5}$$

By applying g_5 to (4.5), we obtain

$$(g_5g_2g_1)(\mathcal{S}_3) + \mathcal{S}_3 \equiv_{\omega_{(4)}} \gamma_{181}a_{179} \equiv_{\omega_{(4)}} 0. \tag{4.6}$$

We have $a_{162} \equiv_{\omega_{(4)}} a_{179} + (g_3g_2g_5g_2g_3)(a_{179})$ and $a_{162} \equiv_{\omega_{(4)}} g_2(a_{161})$. Hence, Lemma 4.3.13 implies that $[a_{179}]_{\omega_{(4)}} \neq 0$. So, from the relation (4.6), we obtain $\gamma_{181} = 0$.

By a simple computation, we can see that the action of Σ_5 on $QP_5(\omega_{(4)})$ induces the one of it on the set $\{[a_t]_{\omega_{(4)}} : 171 \leqslant t \leqslant 200\}$. Furthermore, this action is transitive. Hence, from the relations $\sigma(\mathcal{S}) \equiv_{\omega_{(4)}} 0$ with $\sigma \in \Sigma_5$, we get $\gamma_t = 0$ for $171 \leqslant t \leqslant 200$.

Now, using the above equalities, one gets $\mathcal{S}_2 \equiv_{\omega_{(4)}} \gamma_{208}a_{185} \equiv_{\omega_{(4)}} 0$. This implies $\gamma_{208} = 0$. By using this and the relations $\sigma(\mathcal{S}) \equiv_{\omega_{(4)}} 0$ with $\sigma \in \Sigma_5$, we obtain $\gamma_t = 0$ for $201 \leqslant t \leqslant 210$.

By computing $(g_5g_2g_1)(\mathcal{S}_1) + \mathcal{S}_1$, we have

$$(g_5g_2g_1)(\mathcal{S}_1) + \mathcal{S}_1 \equiv_{\omega_{(4)}} \gamma_{165}a_{162} \equiv_{\omega_{(4)}} 0.$$

Since $[a_{162}]_{\omega_{(4)}} = [g_2(a_{161})]_{\omega_{(4)}} \neq 0$, we get $\gamma_{165} = 0$.

We observe that the action of Σ_5 on $QP_5(\omega_{(4)})$ induces the one on the set $\{[a_t]_{\omega_{(4)}} : 161 \leqslant t \leqslant 170\}$. Since this action is transitive, we get $\gamma_t = 0$ for $161 \leqslant t \leqslant 170$. The proposition is proved. □

Consider the monomials $a_t = a_{35,t},\ 211 \leqslant t \leqslant 225$ as given in Subsection 6.5.

Proposition 4.3.14. *The $\mathbb{F}_2$-vector space $QP_5(\omega_{(5)})$ is an GL_5-module generated by the class $[a_{225}]_{\omega_{(5)}}$ and $B_5(\omega_{(5)}) = \{a_t : 211 \leqslant t \leqslant 225\}$. Consequently,* $\dim QP_5(\omega_{(5)}) = 15$.

We prepare some lemmas for the proof of this proposition. The proof of the following lemma is straightforward.

Lemma 4.3.15. *The following monomials are strictly inadmissible:*

$$x_j^2x_\ell x_t^2x_u^3x_v^3, i < j; \quad x_j^2x_\ell^3x_t^3x_u^3.$$

Here (j, ℓ, t, u, v) is a permutation of $(1, 2, 3, 4, 5)$.

Lemma 4.3.16. *All permutations of the following monomials are strictly inadmissible:*

$$x_1x_2^6x_3^6x_4^7x_5^{15},\ x_1x_2^6x_3^7x_4^7x_5^{14},\ x_1^7x_2^7x_3^9x_4^6x_5^6.$$

Proof. We prove the lemma for $x = x_1x_2^6x_3^6x_4^7x_5^{15}$. By a direct computation, we have

$$x = Sq^1\big(x_1x_2^5x_3^6x_4^7x_5^{15} + x_1^4x_2^3x_3^5x_4^7x_5^{15}\big) + Sq^2\big(x_1^2x_2^3x_3^6x_4^7x_5^{15}\big),\ \ \mathrm{mod}\big(P_5^-(\omega_{(5)})\big).$$

So, all permutations of x are strictly inadmissible. □

Lemma 4.3.17. *The following monomials are strictly inadmissible:*

$$x_1^3x_2^5x_3^6x_4^{14}x_5^7,\ x_1^3x_2^5x_3^{14}x_4^6x_5^7,\ x_1^3x_2^5x_3^{14}x_4^7x_5^6,$$
$$x_1^3x_2^{13}x_3^6x_4^6x_5^7,\ x_1^3x_2^{13}x_3^6x_4^7x_5^6,\ x_1^3x_2^{13}x_3^7x_4^6x_5^6.$$

Proof. We prove the lemma for $x = x_1^3x_2^5x_3^{14}x_4^6x_5^7$. By a direct computation, we have

$$\begin{aligned}x = {} & x_1^3x_2^5x_3^7x_4^6x_5^{14} + Sq^1\big(x_1^3x_2^6x_3^7x_4^5x_5^{13} + x_1^3x_2^6x_3^{13}x_4^5x_5^7\big) + Sq^2\big(x_1^5x_2^3x_3^9x_4^3x_5^{13}\\ & + x_1^5x_2^3x_3^{13}x_4^3x_5^9 + x_1^5x_2^5x_3^7x_4^5x_5^{11} + x_1^5x_2^5x_3^7x_4^9x_5^7 + x_1^5x_2^5x_3^{11}x_4^5x_5^7\\ & + x_1^5x_2^9x_3^7x_4^5x_5^7\big) + Sq^4\big(x_1^3x_2^3x_3^9x_4^3x_5^{13} + x_1^3x_2^3x_3^{13}x_4^3x_5^9 + x_1^3x_2^5x_3^7x_4^5x_5^{11}\\ & + x_1^3x_2^5x_3^7x_4^9x_5^7 + x_1^3x_2^5x_3^9x_4^5x_5^9 + x_1^3x_2^5x_3^{11}x_4^5x_5^7 + x_1^3x_2^9x_3^7x_4^5x_5^7\big)\\ & + Sq^8\big(x_1^3x_2^3x_3^9x_4^3x_5^9\big), \mod\big(P_5^-(\omega_{(5)})\big).\end{aligned}$$

This equality shows that the monomial x is strictly inadmissible. □

Lemma 4.3.18. *The class $[a_{215}]_{\omega_{(5)}}$ is non-zero in the vector space $QP_5(\omega_{(5)})$.*

Proof. Suppose the contrary that $[a_{215}]_{\omega_{(5)}} = 0$. Then

$$a_{215} = \sum_{u=0}^{4} Sq^{2^u}(C_u) \mod(P_5^-(\omega_{(5)})), \tag{4.7}$$

where C_u are suitable polynomials in P_5. Let $(Sq^2)^3$ acts on the both sides of (4.7). Observe that if x is a monomial in $P_5^-(\omega_{(5)})$ then, $(Sq^2)^3(x) \in P_5^-(1,4,4,2)$. Since $(Sq^2)^3Sq^1 = 0$ and $(Sq^2)^3Sq^2 = 0$, we get

$$(Sq^2)^3(a_{215}) = \sum_{u=2}^{4}(Sq^2)^3(Sq^{2^u}(C_u)) \mod(P_5^-(1,4,4,2)).$$

We denote

$$\begin{aligned}&g_{1,2} = x_1^{15}x_2^3x_3^3x_4^5x_5^5,\ g_{1,3} = x_1^{15}x_2^3x_3^5x_4^3x_5^5,\ g_{1,4} = x_1^{15}x_2^3x_3^5x_4^5x_5^3,\\ &g_{2,3} = x_1^{15}x_2^5x_3^3x_4^3x_5^5,\ g_{2,4} = x_1^{15}x_2^5x_3^3x_4^5x_5^3,\ g_{3,4} = x_1^{15}x_2^5x_3^5x_4^3x_5^3.\\ &q_1 = x_1^{15}x_2^3x_3^3x_4^3x_5^9,\ q_2 = x_1^{15}x_2^3x_3^3x_4^9x_5^3,\ q_3 = x_1^{15}x_2^3x_3^9x_4^3x_5^3,\ q_4 = x_1^{15}x_2^9x_3^3x_4^3x_5^3,\\ &r_1 = x_1^{15}x_2^3x_3^3x_4^3x_5^7,\ r_2 = x_1^{15}x_2^3x_3^3x_4^7x_5^3,\ r_3 = x_1^{15}x_2^3x_3^7x_4^3x_5^3,\ r_t = x_1^{15}x_2^7x_3^3x_4^3x_5^3,\\ &s_1 = x_1^{15}x_2^6x_3^6x_4^6x_5^8,\ s_2 = x_1^{15}x_2^6x_3^6x_4^8x_5^6,\ s_3 = x_1^{15}x_2^6x_3^8x_4^6x_5^6,\ s_4 = x_1^{15}x_2^8x_3^6x_4^6x_5^6.\end{aligned}$$

Observe that the polynomial $s = \sum_{u=1}^{4} s_u$ is a term of $(Sq^2)^3(a_{215})$. It is not a term of $(P_5^-(1,4,4,2))$, $(Sq^2)^3(Sq^8(C_3))$ and $(Sq^2)^3(Sq^{16}(C_4))$ hence, it is a term of $(Sq^2)^3(Sq^4(C_2))$. Then, there is (i,j) such that either $g_{i,j}$ or $r = \sum_{u=1}^{4} r_u$ is a term of C_2. If r is a term of C_2, then r_1 is a term of C_2. Then, the monomial $\bar{x} = x_1^{15}x_2^3x_3^3x_4^3x_5^{11}$ is a term of $a_{215} + Sq^4(r_1)$. So $\bar{x}$ is a term of $Sq^4(C_2 + r_1)$. So, r_1 is a term of $C_2 + r_1$. This contradicts the fact that r_1 is a term of C_2.

Thus, we have proved that r is not a term of C_2, hence $g_{i,j}$ is a term of C_2. We can assume that $g_{1,2}$ is a term of C_2.

Now, $x_1^{15}x_2^3x_3^3x_4^9x_5^3$ and $x_1^{15}x_2^3x_3^3x_4^5x_5^9$ are the terms of $a_{215} + Sq^4(g_{1,2})$. Since $r_u, u = 1,2,3,4$ is not a term of C_2, q_1 and q_2 are the term of C_1. Then, the following monomials are the terms of $a_{215} + Sq^4(g_{1,2}) + Sq^2(q_1 + q_2)$:

$$x_1^{15}x_2^5x_3^3x_4^3x_5^9,\ x_1^{15}x_2^3x_3^5x_4^3x_5^9,\ x_1^{15}x_2^5x_3^3x_4^9x_5^3,\ x_1^{15}x_2^3x_3^5x_4^9x_5^3.$$

Since r_u is not a term of C_2, the monomials $g_{1,3}, g_{1,4}, g_{2,3}, g_{2,4}$ are the terms of C_2. Then, $x_1^{15}x_2^3x_3^9x_4^3x_5^5,\ x_1^{15}x_2^9x_3^3x_4^5x_5^3$ are the terms of

$$a_{215} + Sq^4(g_{1,2} + g_{1,3} + g_{1,4} + g_{2,3} + g_{2,4}) + Sq^2(q_1 + q_2).$$

So, $q_3,\ q_4$ are the terms of C_1. Then, $x_1^{15}x_2^5x_3^9x_4^3x_5^3$ is a term of

$$a_{215} + Sq^4(g_{1,2} + g_{1,3} + g_{1,4} + g_{2,3} + g_{2,4}) + Sq^2(q_1 + q_2 + q_3 + q_4).$$

Hence, $g_{3,4}$ is a term of C_2. Now, the monomial $x_1^{15}x_2^6x_3^6x_4^6x_5^8$ is a term of

$$(Sq^2)^3(a_{215} + Sq^4(g_{1,2} + g_{1,3} + g_{1,4} + g_{2,3} + g_{2,4} + g_{3,4})).$$

So, there is (i, j) such that $g_{i,j}$ is a term of $C_2 + \sum_{(i,j)} g_{i,j}$. This contradicts the fact that $g_{i,j}$ is a term of C_2. The lemma is proved. □

Proof of Proposition 4.3.14. Let x be an admissible monomial in P_5 such that $\omega(x) = \omega_{(5)}$. Then $x = x_jx_\ell x_ty^2$ with $y \in B_5(4, 4, 1)$.

Let $z \in B_5(4, 4, 1)$. By a direct computation using the results in Subsection 4.2, we see that if $x_jx_\ell x_tz^2 \neq a_t,\ 211 \leqslant t \leqslant 225$, then there is a monomial w which is given in one of Lemmas 4.3.15 and 4.3.16 such that $x_jx_\ell x_tz^2 = wz_1^{2^u}$ with suitable monomial $z_1 \in P_5$, and $u = \max\{j \in \mathbb{Z} : \omega_j(w) > 0\}$. By Theorem 2.8, $x_jx_\ell x_tz^2$ is inadmissible. Since $x = x_jx_\ell x_ty^2$ and x is admissible, one gets $x = a_t$ for some $t,\ 211 \leqslant t \leqslant 225$. Hence, $B_5(\omega_{(5)}) \subset \{a_t :\ 211 \leqslant t \leqslant 225\}$.

By a direct computation, we see that there is a direct summand decomposition of the Σ_5-modules:

$$QP_5(\omega_{(5)}) = \langle[\Sigma_5(a_{214})]_{\omega_{(5)}}\rangle \bigoplus \langle[\Sigma_5(a_{225})]_{\omega_{(5)}}\rangle.$$

Since $a_{214} \equiv_{\omega_{(5)}} a_{225} + g_5(a_{225})$, $QP_5(\omega_{(5)})$ is the GL_5-module generated by the class $[a_{225}]_{\omega_{(5)}}$.

We now prove the set $\{[a_t]_{\omega_{(5)}} :\ 211 \leqslant t \leqslant 225\}$ is linearly independent in $QP_5(\omega_{(5)})$. Suppose there is a linear relation

$$\mathcal{S} = \sum_{t=211}^{225} \gamma_t a_t \equiv_{\omega_{(5)}} 0, \tag{4.8}$$

where $\gamma_t \in \mathbb{F}_2$. We prove $\gamma_t = 0,\ \forall t,\ 211 \leqslant t \leqslant 225$.

Set $\mathcal{S}_1 = g_5(\mathcal{S}) + \mathcal{S}$. By a direct computation using (4.8), we have

$$(g_5g_2g_1g_2)(\mathcal{S}_1) + \mathcal{S}_1 \equiv_{\omega_{(5)}} \gamma_{215}a_{214} \equiv_{\omega_{(5)}} 0, \tag{4.9}$$

From Lemma 4.3.18, we have $[a_{214}]_{\omega_{(5)}} = [g_1(a_{215}]_{\omega_{(5)}}) \neq 0$. Hence, the relation (4.9) implies $\gamma_{215} = 0$.

Note that the action of Σ_5 on $QP_5(\omega_{(5)})$ induces the one of it on the set $\{[a_t]_{\omega_{(5)}} :\ 211 \leqslant t \leqslant 225\}$ and this action is transitive. So, we get $\gamma_t = 0$ for $211 \leqslant t \leqslant 225$. Then, we have

$$\mathcal{S}_1 \equiv_{\omega_{(5)}} \gamma_{225}a_{214} + \gamma_{222}a_{219} + \gamma_{223}a_{220} + \gamma_{224}a_{221} \equiv_{\omega_{(5)}} 0.$$

The last equalities implies $\gamma_{225} = 0$. Since the action of Σ_5 on the set $\{[a_t]_{\omega_{(5)}} :\ 216 \leqslant t \leqslant 225\}$ is transitive, we obtain $\gamma_t = 0,\ \forall t,\ 216 \leqslant t \leqslant 225$. The proposition is proved. □

Combining the results in this section, one gets the following.

Theorem 4.3.19. *For s an arbitrary positive integer, we have*

$$\dim(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{5(2^s-1)} = \begin{cases} 46, & \text{if } s = 1, \\ 432, & \text{if } s = 2, \\ 1117 & \text{if } s \geqslant 3. \end{cases}$$

This result confirms the one of Hưng in [10], which was proved by using a computer calculation.

5. Proof of Theorem 1.6

In this section, we prove Theorem 1.6 by using the results in Section 4. From Theorem 1.3, we see that the homomorphism

$$(\widetilde{Sq}^0_*)^{s-3} : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)^{GL_5}_{5(2^s-1)} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)^{GL_5}_{35}$$

is an isomorphism for every $s \geqslant 3$. So, we need only to prove the theorem for $s = 1, 2, 3$.

5.1. The Cases $s = 1, 2$.

For $s = 1$, we have $5(2^s - 1) = 5$.

Proposition 5.1.1. $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)^{GL_5}_5 = 0$.

Proof. Denote by $a_t = a_{5,t}$, $1 \leqslant t \leqslant 46$, the admissible monomials of degree 5 in P_5 (see Subsection 6.1). Suppose $f \in (P_5)_5$ such that $[f] \in (\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)^{GL_5}_5$. Then, $f \equiv \sum_{t=1}^{46} \gamma_t a_t$ with $\gamma_t \in \mathbb{F}_2$ and $g_i(f) + f \equiv 0$ for $i = 1, 2, 3, 4, 5$. By computing directly from the relations $g_i(f) + f \equiv 0$, for $i = 1, 2, 3, 4$, we easily obtain $\gamma_t = \gamma_1$, $1 \leqslant t \leqslant 30$, and $\gamma_t = 0$ for $31 \leqslant t \leqslant 45$. Now, by computing $g_5(f) + f$ in terms of the admissible monomials, we get

$$g_5(f) + f \equiv \gamma_1 a_4 + \gamma_{46} a_{31} + \text{ other terms} \equiv 0.$$

The last equality implies $\gamma_1 = \gamma_{46} = 0$. The proposition follows. □

For $s = 2$, the theorem has been proved in [23]. We have

Proposition 5.1.2 (See [23])**.** $(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)^{GL_5}_{15}$ *is an* $\mathbb{F}_2$*-vector space of dimension* 2 *with a basis consisting of the* 2 *classes represented by the polynomials* p *and* q*, which are determined as in Subsection* 6.7*.*

5.2. The Case $s = 3$.

For $s = 3$, $5(2^s - 1) = 35$. From the results in Section 4, we have

$$(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{35} \cong \operatorname{Ker}(\widetilde{Sq}^0_*)_{(5,15)} \bigoplus (\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{15},$$
$$\operatorname{Ker}(\widetilde{Sq}^0_*)_{(5,15)} = \bigoplus_{1 \leqslant j \leqslant 5} QP_5(\omega_{(j)}).$$

Since $QP_5(\omega_{(2)}) = 0$ and $QP_5(\omega_{(3)}) = 0$, we need only to compute $QP_5(\omega_{(j)})^{GL_5}$ for $j = 1, 4, 5$.

5.2.1. *Computation of* $QP_5(\omega_{(1)})^{GL_5}$.

We prove the following.

Proposition 5.2.1. $QP_5(\omega_{(1)})^{GL_5} = 0$.

By a direct computation from Proposition 4.3.2, there is a direct summand decomposition of the Σ_5-modules: $QP_5(\omega_{(1)}) = (QP_5^0)_{35} \bigoplus QP_5^+(\omega_{(1)})$ and

$$(QP_5^0)_{35} = \bigoplus_{i=1}^{4}\langle[\Sigma_5(u_i)]\rangle \bigoplus \langle[\Sigma_5(u_5,u_6,u_7)]\rangle,$$

where

$$u_1 = x_1x_2^3x_3^{31},\ u_2 = x_1x_2^7x_3^{27},\ u_3 = x_1^3x_2^7x_3^{25},\ u_4 = x_1x_2x_3^2x_4^{31},$$
$$u_5 = x_1x_2^2x_3^3x_4^{29},\ u_6 = x_1x_2^2x_3^5x_4^{27},\ u_7 = x_1x_2^3x_3^5x_4^{26}.$$

We denote the admissible monomials in $(QP_5^0)_{35}$ by $b_t,\ 1 \leqslant t \leqslant 460$, as given in Subsection 6.5. We prepare some lemmas for the proof of the proposition.

Lemma 5.2.2.

i) $\langle[\Sigma_5(u_i)]\rangle^{\Sigma_5} = \langle[p_i]\rangle, i = 1,2,3;\ \langle[\Sigma_5(u_4)]\rangle^{\Sigma_5} = 0$.

ii) $\langle[\Sigma_5(u_5,u_6,u_7)]\rangle^{\Sigma_5} = \langle[p_4],[p_5],[p_6]\rangle$.

Here, the polynomials $p_i,\ 1 \leqslant i \leqslant 6$, *are determined as in Subsection 6.6.*

Outline of the proof. The set $B_1 := \{[b_t] : 1 \leqslant t \leqslant 60\}$ is a basis of $\langle[\Sigma_5(u_1)]\rangle$. The action of Σ_5 on $QP_5(\omega_{(1)})$ induces the one of it on B_1. Furthermore, this action is transitive. Hence, if $f = \sum_{t=1}^{60}\gamma_t b_t$ with $\gamma_t \in \mathbb{F}_2$ and $[f] \in \langle[\Sigma_5(u_1)]\rangle^{\Sigma_5}$, then the relations $g_j(f) \equiv f, j = 1,2,3,4$, imply $\gamma_t = \gamma_1, \forall t,\ 1 \leqslant t \leqslant 60$. The first part of the lemma is proved for $i = 1$. The case $i = 2$ is proved by a similar argument.

The set $B_3 := \{[b_t] : 91 \leqslant t \leqslant 140\}$ is the basis of $\langle[\Sigma_5(u_3)]\rangle$. Suppose $f = \sum_{t=91}^{140}\gamma_t b_t$ with $\gamma_t \in \mathbb{F}_2$ and $[f] \in \langle[\Sigma_5(u_3)]\rangle^{\Sigma_5}$. By computing directly the relations $g_j(f) \equiv f, j = 1,2,3,4$, we get $\gamma_t = 0$ for $t \in \mathbb{J}_1$ and $\gamma_t = \gamma_{92}$ for $t \notin \mathbb{J}_1$, where $\mathbb{J}_1 = \{91,93,95,96,99,101,102,105,106,107\}$. So, $f \equiv p_3$.

The set $B_4 := \{[b_t] : 141 \leqslant t \leqslant 180\}$ is the basis of $\langle[\Sigma_5(u_4)]\rangle$. Suppose $f = \sum_{t=141}^{180}\gamma_t b_t$ with $\gamma_t \in \mathbb{F}_2$ and $[f] \in \langle[\Sigma_5(u_4)]\rangle^{\Sigma_5}$. By computing from the relations $g_j(f) \equiv f, j = 1,2,3,4$, we obtain $\gamma_t = 0, \forall t, 141 \leqslant t \leqslant 180$. Hence, $f \equiv 0$.

The set $B_5 := \{[b_t] : 181 \leqslant t \leqslant 460\}$ is the basis of $\langle[\Sigma_5(u_5,u_6,u_7)]\rangle$. Suppose $f = \sum_{t=141}^{180}\gamma_t b_t$ with $\gamma_t \in \mathbb{F}_2$ and $[f] \in \langle[\Sigma_5(u_5,u_6,u_7)]\rangle^{\Sigma_5}$. By a direct computation from the relations $g_j(f) \equiv f, j = 1,2,3,4$, we get $f \equiv \gamma_{181}p_4 + \gamma_{189}p_5 + \gamma_{198}p_6$. The second part is proved. □

Lemma 5.2.3. $QP_5^+(\omega_{(1)})^{\Sigma_5} = \langle[p_7],[p_8],[p_9]\rangle$. *Here, the polynomials* $p_i,\ 7 \leqslant i \leqslant 9$, *are determined as in Subsection 6.6.*

Outline of the proof. Let $[f] \in QP_5^+(\omega_{(1)})^{\Sigma_5}$. By Proposition 4.3.2, $f \equiv \sum_{t=1}^{160}\gamma_t a_t$. We compute $g_j(f) + f$ in terms of $a_t, 1 \leqslant t \leqslant 160$, $(\mathrm{mod}(\mathcal{A}^+P_5)$. By computing from the relations $g_j(f) + f \equiv 0$ with $j = 1,2,3,4$, we get $f \equiv \gamma_1p_7 + \gamma_4p_8 + \gamma_{13}p_9$. Hence, the lemma follows. □

Proof of Proposition 5.2.1. Combining Lemmas 5.2.2 and 5.2.3 gives

$$QP_5(\omega_{(1)})^{\Sigma_5} = \langle\{[p_j] : 1 \leqslant j \leqslant 9\}\rangle.$$

Let $f \in P_5(\omega_{(1)})$ such that $[f] \in QP_5(\omega_{(1)})^{GL_5}$. Then, $f \equiv \sum_{j=1}^{9} \gamma_j p_j$ with $\gamma_j \in \mathbb{F}_2$. By a direct computation using Theorem 2.11, we have

$$\begin{aligned} g_5(f) + f \equiv{}& \gamma_1 b_7 + \gamma_2 b_{48} + \gamma_3 b_{36} + \gamma_4 b_{172} + \gamma_5 b_{398} + (\gamma_1 + \gamma_5 + \gamma_6) b_{21} \\ &+ \gamma_7 b_{371} + \gamma_8 a_{136} + \gamma_9 b_{182} + \text{ other terms} \equiv 0. \end{aligned}$$

The last equality implies $\gamma_t = 0$ for $1 \leqslant t \leqslant 9$. The proposition is proved. □

5.2.2. *Computation of* $QP_5(\omega_{(4)})^{GL_5}$.

In this part, we prove the following.

Proposition 5.2.4. $QP_5(\omega_{(4)})^{GL_5} = 0$.

From Proposition 4.3.10, we can see that there is a direct summand decomposition of the Σ_5-modules:

$$QP_5(\omega_{(4)}) = \langle [\Sigma_5(a_{162})]_{\omega_{(4)}} \rangle \bigoplus \langle [\Sigma_5(a_{175})]_{\omega_{(4)}} \rangle \bigoplus \langle [\Sigma_5(a_{203})]_{\omega_{(4)}} \rangle.$$

Lemma 5.2.5. $\langle [\Sigma_5(a_{162})]_{\omega_{(4)}} \rangle^{\Sigma_5} = \langle [p(a_{162})]_{\omega_{(4)}} \rangle$.

Proof. The set $\{[a_t]_{\omega_{(4)}} : 161 \leqslant t \leqslant 170\}$ is a basis of $\langle [\Sigma_5(a_{162})]_{\omega_{(4)}} \rangle$. If $f \in P_5(\omega_{(4)})$ such that $[f]_{\omega_{(4)}} \in \langle [\Sigma_5(a_{162})]_{\omega_{(4)}} \rangle^{\Sigma_5}$, then $f \equiv_{\omega_{(4)}} \sum_{t=161}^{170} \gamma_t a_t$ with $\gamma_t \in \mathbb{F}_2$ and $g_j(f) + f \equiv_{\omega_{(4)}} 0$, for $j = 1, 2, 3, 4$. Since the action of Σ_5 on $QP_5(\omega_{(4)})$ induces the one of it on $\{[a_t]_{\omega_{(4)}} : 161 \leqslant t \leqslant 170\}$ which is transitive. Hence, from the relation $g_j(f) + f \equiv_{\omega_{(4)}} 0$, with $j = 1, 2, 3, 4$, we obtain $\gamma_t = \gamma_{161}, \forall t, 161 \leqslant t \leqslant 170$. The lemma follows. □

By an argument similar to the proof of Lemma 5.2.5, we get the following.

Lemma 5.2.6. $\langle [\Sigma_5(a_{175})]_{\omega_{(4)}} \rangle^{\Sigma_5} = \langle [p(a_{175})]_{\omega_{(4)}} \rangle$.

Lemma 5.2.7. $\langle [\Sigma_5(a_{203})]_{\omega_{(4)}} \rangle^{\Sigma_5} = 0$.

Proof. The set $\{[a_t]_{\omega_{(4)}} : 201 \leqslant t \leqslant 210\}$ is a basis of $\langle [\Sigma_5(a_{203})]_{\omega_{(4)}} \rangle$. If $f \in P_5(\omega_{(4)})$ such that $[f]_{\omega_{(4)}} \in \langle [\Sigma_5(a_{204})]_{\omega_{(4)}} \rangle^{\Sigma_5}$, then $f \equiv_{\omega_{(4)}} \sum_{t=201}^{210} \gamma_t a_t$ with $\gamma_t \in \mathbb{F}_2$ and $g_j(f) + f \equiv_{\omega_{(4)}} 0$, for $j = 1, 2, 3, 4$. By a direct computation, we get

$$\begin{aligned} g_1(f) + f \equiv_{\omega_{(4)}}{}& \gamma_{204} a_{201} + \gamma_{205} a_{202} + \gamma_{206} a_{203} + (\gamma_{207} + \gamma_{209}) a_{207} \\ &+ (\gamma_{208} + \gamma_{210}) a_{208} + (\gamma_{207} + \gamma_{209}) a_{209} + (\gamma_{208} + \gamma_{210}) a_{210} \equiv_{\omega_{(4)}} 0, \\ g_2(f) + f \equiv_{\omega_{(4)}}{}& (\gamma_{201} + \gamma_{204}) a_{201} + (\gamma_{202} + \gamma_{205}) a_{202} + (\gamma_{203} + \gamma_{207} + \gamma_{208}) a_{203} \\ &+ (\gamma_{201} + \gamma_{204}) a_{204} + (\gamma_{202} + \gamma_{205}) a_{205} + (\gamma_{206} + \gamma_{208}) a_{206} \\ &+ (\gamma_{203} + \gamma_{206} + \gamma_{207}) a_{207} + (\gamma_{206} + \gamma_{208}) a_{208} + \gamma_{210} a_{209} \equiv_{\omega_{(4)}} 0, \\ g_3(f) + f \equiv_{\omega_{(4)}}{}& \gamma_{204} a_{201} + (\gamma_{202} + \gamma_{203} + \gamma_{206}) a_{202} + (\gamma_{202} + \gamma_{203} + \gamma_{205}) a_{203} \\ &+ (\gamma_{205} + \gamma_{206}) a_{205} + (\gamma_{205} + \gamma_{206}) a_{206} + (\gamma_{207} + \gamma_{208}) a_{207} \\ &+ (\gamma_{207} + \gamma_{208}) a_{208} + (\gamma_{209} + \gamma_{210}) a_{209} + (\gamma_{209} + \gamma_{210}) a_{210} \equiv_{\omega_{(4)}} 0, \\ g_4(f) + f \equiv_{\omega_{(4)}}{}& (\gamma_{201} + \gamma_{202}) a_{201} + (\gamma_{201} + \gamma_{202}) a_{202} + \gamma_{206} a_{203} \\ &+ (\gamma_{204} + \gamma_{205}) a_{204} + (\gamma_{204} + \gamma_{205}) a_{205} + \gamma_{208} a_{207} + \gamma_{210} a_{209} \equiv_{\omega_{(4)}} 0. \end{aligned}$$

From the above equalities, we obtain $\gamma_t = 0, \forall t, 201 \leqslant t \leqslant 210$. The lemma is proved. □

Proof of Proposition 5.2.4. By combining Lemmas 5.2.5-5.2.7, we have

$$QP_5(\omega_{(4)})^{\Sigma_5} = \langle [p(a_{162})]_{\omega_{(4)}}, [p(a_{174})]_{\omega_{(4)}} \rangle,$$

where $p(a_{162}) = \sum_{t=161}^{170} a_t$ and $p(a_{174}) = \sum_{t=171}^{200} a_t$.

Let $f \in P_5(\omega_{(4)})$ such that $[f] \in QP_5(\omega_{(4)})^{GL_5}$. Then, $f \equiv_{\omega_{(4)}} \gamma_1 p(a_{162}) + \gamma_2 p(a_{174})$ with $\gamma_1, \gamma_2 \in \mathbb{F}_2$. By a direct computation, we have

$$g_5(f) + f \equiv_{\omega_{(4)}} (\gamma_1 + \gamma_2)a_{162} + \gamma_2 a_{171} + \text{other terms} \equiv_{\omega_{(4)}} 0.$$

This equality implies $\gamma_1 = \gamma_2 = 0$. The proposition is proved. □

5.2.3. *Computation of* $QP_5(\omega_{(5)})^{GL_5}$.

In this part, we prove the following.

Proposition 5.2.8. $QP_5(\omega_{(5)})^{GL_5} = 0$.

From Proposition 4.3.14, there is a direct summand decomposition of the Σ_5-modules:

$$QP_5(\omega_{(5)}) = \langle [\Sigma_5(a_{215})]_{\omega_{(5)}} \rangle \bigoplus \langle [\Sigma_5(a_{225})]_{\omega_{(5)}} \rangle.$$

By an argument similar to the proof of Lemma 5.2.5, we get the following.

Lemma 5.2.9. $\langle [\Sigma_5(a_i)]_{\omega_{(5)}} \rangle^{\Sigma_5} = \langle [p(a_i)]_{\omega_{(5)}} \rangle$ *with* $i = 215, 225$.

Proof of Proposition 5.2.8. From Lemma 5.2.9, we have

$$QP_5(\omega_{(5)})^{\Sigma_5} = \langle [p(a_{215})]_{\omega_{(5)}}, [p(a_{225})]_{\omega_{(5)}} \rangle,$$

where $p(a_{215}) = \sum_{t=211}^{215} a_t$ and $p(a_{225}) = \sum_{t=216}^{225} a_t$.

Let $f \in P_5(\omega_{(5)})$ such that $[f] \in QP_5(\omega_{(5)})^{GL_5}$. Then, $f \equiv_{\omega_{(5)}} \gamma_1 p(a_{215}) + \gamma_2 p(a_{225})$ with $\gamma_1, \gamma_2 \in \mathbb{F}_2$. A direct computation shows

$$g_5(f) + f \equiv_{\omega_{(5)}} (\gamma_1 + \gamma_2)a_{214} + \gamma_2(a_{219} + a_{220} + a_{221}) \equiv_{\omega_{(5)}} 0.$$

This equality implies $\gamma_1 = \gamma_2 = 0$. The proposition is proved. □

From the above results, we easily obtain the following.

Corollary 5.2.10. $\mathrm{Ker}(\widetilde{Sq}^0_*)^{GL_5}_{(5,15)} = 0$.

5.2.4. *Proof of Theorem 1.6 for* $s = 3$.

Let $f \in (P_5)_{35}$ such that $[f] \in (\mathbb{F}_2 \otimes P_5)^{GL_5}_{35}$. Since Kameko's squaring operation

$$(\widetilde{Sq}^0_*)_{(5,15)} : (\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{35} \longrightarrow (\mathbb{F}_2 \otimes_{\mathcal{A}} P_k)_{15}$$

is an epimorphism of GL_5-modules, $(\widetilde{Sq}^0_*)_{(5,15)}([f]) \in (\mathbb{F}_2 \otimes P_5)^{GL_5}_{15}$. By Proposition 5.1.2, $(\widetilde{Sq}^0_*)_{(5,15)}([f]) = \lambda_1[p] + \lambda_2[q]$ with $\lambda_1, \lambda_2 \in \mathbb{F}_2$. Hence, we have

$$f \equiv \lambda_1\psi(p) + \lambda_2\psi(q) + \bar{f},$$

where $\bar{f} \in (P_5)_{35}$ such that $[\bar{f}] \in \mathrm{Ker}(\widetilde{Sq}^0_*)_{(5,15)}$, and $\psi : P_5 \to P_5$ is the $\mathbb{F}_2$-linear map determined by $\psi(y) = x_1x_2x_3x_4x_5y^2$ for any $y \in P_5$.

Now, we prove that if $[f] \neq 0$, then $\lambda_2 = 1$.

Suppose the contrary, that $\lambda_2 = 0$. By a direct computation, we have

$$g_1(\psi(p)) + \psi(p) \equiv 0, \tag{5.1}$$

$$g_2(\psi(p)) + \psi(p) \equiv a_{45} + a_{46} + a_{49} + a_{50} + a_{53} + a_{54} + a_{61} + a_{62} + a_{72} + a_{73} + a_{76} + a_{77} + a_{80} + a_{81} + a_{91} + a_{92}, \tag{5.2}$$

$$g_3(\psi(p)) + \psi(p) \equiv a_8 + a_{12} + a_{15} + a_{17} + a_{36} + a_{38} + a_{40} + a_{49} + a_{53} + a_{60}, \tag{5.3}$$

$$g_4(\psi(p)) + \psi(p) \equiv a_2 + a_6 + a_8 + a_9 + a_{11} + a_{27} + a_{29} + a_{30} + a_{32} + a_{40} + a_{43}. \tag{5.4}$$

From the relations (5.1)-(5.4), we have $g_i(\bar{f}) + \bar{f} \equiv_{\omega_{(t)}} g_i(f) + f \equiv_{\omega_{(t)}} 0$, for $i = 1, 2, 3, 4, 5$ and $t = 4, 5$. From this and Propositions 5.2.4 and 5.2.8, we get $[\bar{f}]_{\omega_{(t)}} \in QP_5(\omega_{(t)})^{GL_5} = 0$. Combining this and the facts that $QP_5(\omega_{(2)}) = 0$ and $QP_5(\omega_{(3)}) = 0$ gives $\bar{f} \equiv f' \in QP_5(\omega_{(1)})$. Now, by a direct computation from the relations $g_i(f) + f \equiv 0$, for $i = 1, 2, 3, 4$, and using Proposition 4.3.2, we get $\lambda_1 = 0$. Hence, $[f] = [f'] \in QP_5(\omega_{(1)})^{GL_5}$. By Proposition 5.2.1, $[f] = 0$. This contradicts the hypothesis $[f] \neq 0$. Hence, $\lambda_2 = 1$ and $f \equiv \lambda_1\psi(p) + \psi(q) + \bar{f}$.

Suppose that $f^* = \lambda\psi(p) + \psi(q) + \bar{f}^*$ with $\lambda \in \mathbb{F}_2$, $[\bar{f}^*] \in \mathrm{Ker}(\widetilde{Sq}^0_*)_{(5,15)}$ and $[f^*] \in (\mathbb{F}_2 \otimes P_5)^{GL_5}_{35}$. Then, $(\lambda_1 + \lambda)\psi(p) + \bar{f} + \bar{f}^* = f + f^*$ and

$$[f + f^*] = [f] + [f^*] \in (\mathbb{F}_2 \otimes P_5)^{GL_5}_{35}.$$

Hence, $\lambda = \lambda_1$ and $[\bar{f} + \bar{f}^*] \in \mathrm{Ker}(\widetilde{Sq}^0_*)^{GL_5}_{(5,15)} = 0$. This implies $[f] = [f^*]$.

Thus, we have proved that $\dim(\mathbb{F}_2 \otimes P_5)^{GL_5}_{35} \leqslant 1$.

In [19], Singer showed that the Adams elements h_i are in the image of φ_1^*. Hà showed in [8] that the elements d_i are in the image of φ_4^*. Since $\varphi^* = \bigoplus_{k\geqslant 0}\varphi_k^*$ is a homomorphism of algebras, we see that the element h_2d_1 is in the image of φ_5^*, hence $\varphi_5((h_2d_1)^*) \neq 0$. This implies $\dim(\mathbb{F}_2 \otimes P_5)^{GL_5}_{35} \geqslant 1$. Theorem 1.6 is completely proved.

6. Appendix

In the appendix, we list all admissible monomials of degrees 5, 7, 15, 16, 35 in P_4 and P_5. We order a set of some monomials in P_k by using the order as in Definition 2.5.

6.1. The admissible monomials of degree 5 in $\mathbf{P_5}$.

$B_5(5)$ is the set of 46 monomials:

1. $x_3x_4x_5^3$	2. $x_3x_4^3x_5$	3. $x_3^3x_4x_5$	4. $x_2x_4x_5^3$
5. $x_2x_4^3x_5$	6. $x_2x_3x_5^3$	7. $x_2x_3x_4^3$	8. $x_2x_3^3x_5$
9. $x_2x_3^3x_4$	10. $x_2^3x_4x_5$	11. $x_2^3x_3x_5$	12. $x_2^3x_3x_4$
13. $x_1x_4x_5^3$	14. $x_1x_4^3x_5$	15. $x_1x_3x_5^3$	16. $x_1x_3x_4^3$
17. $x_1x_3^3x_5$	18. $x_1x_3^3x_4$	19. $x_1x_2x_5^3$	20. $x_1x_2x_4^3$
21. $x_1x_2x_3^3$	22. $x_1x_2^3x_5$	23. $x_1x_2^3x_4$	24. $x_1x_2^3x_3$
25. $x_1^3x_4x_5$	26. $x_1^3x_3x_5$	27. $x_1^3x_3x_4$	28. $x_1^3x_2x_5$
29. $x_1^3x_2x_4$	30. $x_1^3x_2x_3$	31. $x_2x_3x_4x_5^2$	32. $x_2x_3x_4^2x_5$

33.	$x_2x_3^2x_4x_5$	34.	$x_1x_3x_4x_5^2$	35.	$x_1x_3x_4^2x_5$	36.	$x_1x_3^2x_4x_5$
37.	$x_1x_2x_4x_5^2$	38.	$x_1x_2x_4^2x_5$	39.	$x_1x_2x_3x_5^2$	40.	$x_1x_2x_3x_4^2$
41.	$x_1x_2x_3^2x_5$	42.	$x_1x_2x_3^2x_4$	43.	$x_1x_2^2x_4x_5$	44.	$x_1x_2^2x_3x_5$
45.	$x_1x_2^2x_3x_4$	46.	$x_1x_2x_3x_4x_5$.				

6.2. **The admissible monomials of degree 7 in $\mathbf{P_5}$.**

6.2.1. $B_4(7)$ is the set of 35 monomials:

1. x_4^7	2. $x_3x_4^6$	3. x_3^7	4. $x_2x_4^6$	5. $x_2x_3^2x_4^4$
6. $x_2x_3^6$	7. x_2^7	8. $x_1x_4^6$	9. $x_1x_3^2x_4^4$	10. $x_1x_3^6$
11. $x_1x_2^2x_4^4$	12. $x_1x_2^2x_3^4$	13. $x_1x_2^6$	14. x_1^7	15. $x_1x_2^2x_3^2x_4^2$
16. $x_2x_3^3x_4^3$	17. $x_2^3x_3x_4^3$	18. $x_2^3x_3^3x_4$	19. $x_1x_3^3x_4^3$	20. $x_1x_2x_3^2x_4^3$
21. $x_1x_2x_3^3x_4^2$	22. $x_1x_2^2x_3x_4^3$	23. $x_1x_2^2x_3^3x_4$	24. $x_1x_2^3x_4^3$	25. $x_1x_2^3x_3x_4^2$
26. $x_1x_2^3x_3^2x_4$	27. $x_1x_2^3x_3^3$	28. $x_1^3x_3x_4^3$	29. $x_1^3x_3^3x_4$	30. $x_1^3x_2x_4^3$
31. $x_1^3x_2x_3x_4^2$	32. $x_1^3x_2x_3^2x_4$	33. $x_1^3x_2x_3^3$	34. $x_1^3x_2^3x_4$	35. $x_1^3x_2^3x_3$

6.2.2. $B_5(7) = f(B_4(7)) \cup B_5^+(3,2) \cup B_5(5,1)$, where

$B_5^+(3,2) = \{x_1x_2x_3x_4^2x_5^2,\ x_1x_2x_3^2x_4x_5^2,\ x_1x_2x_3^2x_4^2x_5,\ x_1x_2^2x_3x_4x_5^2,\ x_1x_2^2x_3x_4^2x_5\}$,
$B_5^+(5,1) = \{x_1x_2x_3x_4x_5^3,\ x_1x_2x_3x_4^3x_5,\ x_1x_2x_3^3x_4x_5,\ x_1x_2^3x_3x_4x_5,\ x_1^3x_2x_3x_4x_5\}$.

We have $|f(B_4(7))| = 100$. Hence, $\dim(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_7 = 110$.

6.3. **The admissible monomials of degree 15 in $\mathbf{P_4}$.**

6.3.1. $B_4(15)$ is the set of 75 monomials:

1. x_4^{15}	2. $x_3x_4^{14}$	3. x_3^{15}	4. $x_2x_4^{14}$	5. $x_2x_3^2x_4^{12}$
6. $x_2x_3^{14}$	7. x_2^{15}	8. $x_1x_4^{14}$	9. $x_1x_3^2x_4^{12}$	10. $x_1x_3^{14}$
11. $x_1x_2^2x_4^{12}$	12. $x_1x_2^2x_3^4x_4^8$	13. $x_1x_2^2x_3^{12}$	14. $x_1x_2^{14}$	15. x_1^{15}
16. $x_2x_3^7x_4^7$	17. $x_2^3x_3^5x_4^7$	18. $x_2^3x_3^7x_4^5$	19. $x_2^7x_3x_4^7$	20. $x_2^7x_3^3x_4^5$
21. $x_2^7x_3^7x_4$	22. $x_1x_3^7x_4^7$	23. $x_1x_2x_3^6x_4^7$	24. $x_1x_2x_3^7x_4^6$	25. $x_1x_2^2x_3^5x_4^7$
26. $x_1x_2^2x_3^7x_4^5$	27. $x_1x_2^3x_3^4x_4^7$	28. $x_1x_2^3x_3^5x_4^6$	29. $x_1x_2^3x_3^6x_4^5$	30. $x_1x_2^3x_3^7x_4^4$
31. $x_1x_2^6x_3x_4^7$	32. $x_1x_2^6x_3^3x_4^5$	33. $x_1x_2^6x_3^7x_4$	34. $x_1x_2^7x_4^7$	35. $x_1x_2^7x_3x_4^6$
36. $x_1x_2^7x_3^2x_4^5$	37. $x_1x_2^7x_3^3x_4^4$	38. $x_1x_2^7x_3^6x_4$	39. $x_1x_2^7x_3^7$	40. $x_1^3x_3^5x_4^7$
41. $x_1^3x_3^7x_4^5$	42. $x_1^3x_2x_3^4x_4^7$	43. $x_1^3x_2x_3^5x_4^6$	44. $x_1^3x_2x_3^6x_4^5$	45. $x_1^3x_2x_3^7x_4^4$
46. $x_1^3x_2^3x_3^4x_4^5$	47. $x_1^3x_2^3x_3^5x_4^4$	48. $x_1^3x_2^4x_3x_4^7$	49. $x_1^3x_2^4x_3^3x_4^5$	50. $x_1^3x_2^4x_3^7x_4$
51. $x_1^3x_2^5x_4^7$	52. $x_1^3x_2^5x_3x_4^6$	53. $x_1^3x_2^5x_3^2x_4^5$	54. $x_1^3x_2^5x_3^3x_4^4$	55. $x_1^3x_2^5x_3^6x_4$
56. $x_1^3x_2^5x_3^7$	57. $x_1^3x_2^7x_4^5$	58. $x_1^3x_2^7x_3x_4^4$	59. $x_1^3x_2^7x_3^4x_4$	60. $x_1^3x_2^7x_3^5$
61. $x_1^7x_3x_4^7$	62. $x_1^7x_3^3x_4^5$	63. $x_1^7x_3^7x_4$	64. $x_1^7x_2x_4^7$	65. $x_1^7x_2x_3x_4^6$
66. $x_1^7x_2x_3^2x_4^5$	67. $x_1^7x_2x_3^3x_4^4$	68. $x_1^7x_2x_3^6x_4$	69. $x_1^7x_2x_3^7$	70. $x_1^7x_2^3x_4^5$
71. $x_1^7x_2^3x_3x_4^4$	72. $x_1^7x_2^3x_3^4x_4$	73. $x_1^7x_2^3x_3^5$	74. $x_1^7x_2^7x_4$	75. $x_1^7x_2^7x_3$

$B_5(15) = f(B_4(15)) \cup B_5(1,1,3) \cup B_5^+(3,2,2) \cup B_5(3,4,1) \cup \psi(B_5(5)$, where

6.3.2. $B_5(1,1,3) = \{x_1x_2^2x_3^4x_4^4x_5^4\}$;

6.3.3. $B_5^+(3,2,2)$ is the set of 75 monomials:

1. $x_1x_2x_3x_4^6x_5^6$	2. $x_1x_2x_3^2x_4^4x_5^7$	3. $x_1x_2x_3^2x_4^5x_5^6$	4. $x_1x_2x_3^2x_4^6x_5^5$
5. $x_1x_2x_3^2x_4^7x_5^4$	6. $x_1x_2x_3^3x_4^4x_5^6$	7. $x_1x_2x_3^3x_4^6x_5^4$	8. $x_1x_2x_3^6x_4x_5^6$
9. $x_1x_2x_3^6x_4^2x_5^5$	10. $x_1x_2x_3^6x_4^3x_5^4$	11. $x_1x_2x_3^6x_4^6x_5$	12. $x_1x_2x_3^7x_4^2x_5^4$
13. $x_1x_2^2x_3x_4^4x_5^7$	14. $x_1x_2^2x_3x_4^5x_5^6$	15. $x_1x_2^2x_3x_4^6x_5^5$	16. $x_1x_2^2x_3x_4^7x_5^4$
17. $x_1x_2^2x_3^3x_4^4x_5^5$	18. $x_1x_2^2x_3^3x_4^5x_5^4$	19. $x_1x_2^2x_3^4x_4x_5^7$	20. $x_1x_2^2x_3^4x_4^3x_5^5$
21. $x_1x_2^2x_3^4x_4^7x_5$	22. $x_1x_2^2x_3^5x_4x_5^6$	23. $x_1x_2^2x_3^5x_4^2x_5^5$	24. $x_1x_2^2x_3^5x_4^3x_5^4$
25. $x_1x_2^2x_3^5x_4^6x_5$	26. $x_1x_2^2x_3^7x_4x_5^4$	27. $x_1x_2^2x_3^7x_4^4x_5$	28. $x_1x_2^3x_3x_4^4x_5^6$
29. $x_1x_2^3x_3x_4^6x_5^4$	30. $x_1x_2^3x_3^2x_4^4x_5^5$	31. $x_1x_2^3x_3^2x_4^5x_5^4$	32. $x_1x_2^3x_3^3x_4^4x_5^4$
33. $x_1x_2^3x_3^4x_4x_5^6$	34. $x_1x_2^3x_3^4x_4^2x_5^5$	35. $x_1x_2^3x_3^4x_4^3x_5^4$	36. $x_1x_2^3x_3^4x_4^6x_5$
37. $x_1x_2^3x_3^5x_4^2x_5^4$	38. $x_1x_2^3x_3^6x_4x_5^4$	39. $x_1x_2^3x_3^6x_4^4x_5$	40. $x_1x_2^6x_3x_4x_5^6$
41. $x_1x_2^6x_3x_4^2x_5^5$	42. $x_1x_2^6x_3x_4^3x_5^4$	43. $x_1x_2^6x_3x_4^6x_5$	44. $x_1x_2^6x_3^3x_4x_5^4$
45. $x_1x_2^6x_3^3x_4^4x_5$	46. $x_1x_2^7x_3x_4^2x_5^4$	47. $x_1x_2^7x_3^2x_4x_5^4$	48. $x_1x_2^7x_3^2x_4^4x_5$
49. $x_1^3x_2x_3x_4^4x_5^6$	50. $x_1^3x_2x_3x_4^6x_5^4$	51. $x_1^3x_2x_3^2x_4^4x_5^5$	52. $x_1^3x_2x_3^2x_4^5x_5^4$
53. $x_1^3x_2x_3^3x_4^4x_5^4$	54. $x_1^3x_2x_3^4x_4x_5^6$	55. $x_1^3x_2x_3^4x_4^2x_5^5$	56. $x_1^3x_2x_3^4x_4^3x_5^4$
57. $x_1^3x_2x_3^4x_4^6x_5$	58. $x_1^3x_2x_3^5x_4^2x_5^4$	59. $x_1^3x_2x_3^6x_4x_5^4$	60. $x_1^3x_2x_3^6x_4^4x_5$
61. $x_1^3x_2^3x_3x_4^4x_5^4$	62. $x_1^3x_2^3x_3^4x_4x_5^4$	63. $x_1^3x_2^3x_3^4x_4^4x_5$	64. $x_1^3x_2^4x_3x_4x_5^6$
65. $x_1^3x_2^4x_3x_4^2x_5^5$	66. $x_1^3x_2^4x_3x_4^3x_5^4$	67. $x_1^3x_2^4x_3x_4^6x_5$	68. $x_1^3x_2^4x_3^3x_4x_5^4$
69. $x_1^3x_2^4x_3^3x_4^4x_5$	70. $x_1^3x_2^5x_3x_4^2x_5^4$	71. $x_1^3x_2^5x_3^2x_4x_5^4$	72. $x_1^3x_2^5x_3^2x_4^4x_5$
73. $x_1^7x_2x_3x_4^2x_5^4$	74. $x_1^7x_2x_3^2x_4x_5^4$	75. $x_1^7x_2x_3^2x_4^4x_5$	

6.3.4. $B_5(3,4,1)$ is the set of 40 monomials:

1. $x_1x_2^2x_3^2x_4^3x_5^7$	2. $x_1x_2^2x_3^2x_4^7x_5^3$	3. $x_1x_2^2x_3^3x_4^2x_5^7$	4. $x_1x_2^2x_3^3x_4^3x_5^6$
5. $x_1x_2^2x_3^3x_4^6x_5^3$	6. $x_1x_2^2x_3^3x_4^7x_5^2$	7. $x_1x_2^2x_3^7x_4^2x_5^3$	8. $x_1x_2^2x_3^7x_4^3x_5^2$
9. $x_1x_2^3x_3^2x_4^2x_5^7$	10. $x_1x_2^3x_3^2x_4^3x_5^6$	11. $x_1x_2^3x_3^2x_4^6x_5^3$	12. $x_1x_2^3x_3^2x_4^7x_5^2$
13. $x_1x_2^3x_3^3x_4^2x_5^6$	14. $x_1x_2^3x_3^3x_4^6x_5^2$	15. $x_1x_2^3x_3^6x_4^2x_5^3$	16. $x_1x_2^3x_3^6x_4^3x_5^2$
17. $x_1x_2^3x_3^7x_4^2x_5^2$	18. $x_1x_2^7x_3^2x_4^2x_5^3$	19. $x_1x_2^7x_3^2x_4^3x_5^2$	20. $x_1x_2^7x_3^3x_4^2x_5^2$
21. $x_1^3x_2x_3^2x_4^2x_5^7$	22. $x_1^3x_2x_3^2x_4^3x_5^6$	23. $x_1^3x_2x_3^2x_4^6x_5^3$	24. $x_1^3x_2x_3^2x_4^7x_5^2$
25. $x_1^3x_2x_3^3x_4^2x_5^6$	26. $x_1^3x_2x_3^3x_4^6x_5^2$	27. $x_1^3x_2x_3^6x_4^2x_5^3$	28. $x_1^3x_2x_3^6x_4^3x_5^2$
29. $x_1^3x_2x_3^7x_4^2x_5^2$	30. $x_1^3x_2^3x_3x_4^2x_5^6$	31. $x_1^3x_2^3x_3x_4^6x_5^2$	32. $x_1^3x_2^3x_3^5x_4^2x_5^2$
33. $x_1^3x_2^5x_3^2x_4^2x_5^3$	34. $x_1^3x_2^5x_3^2x_4^3x_5^2$	35. $x_1^3x_2^5x_3^3x_4^2x_5^2$	36. $x_1^3x_2^7x_3x_4^2x_5^2$
37. $x_1^7x_2x_3^2x_4^2x_5^3$	38. $x_1^7x_2x_3^2x_4^3x_5^2$	39. $x_1^7x_2x_3^3x_4^2x_5^2$	40. $x_1^7x_2^3x_3x_4^2x_5^2$

We have $|f(B_4(15)| = 270$. Hence, $\dim(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)_{15} = 432$.

6.4. **The admissible monomials of degree 16 in $\mathbf{P_5}$.**

6.4.1. $B_4(16)$ is the set of 73 monomials:

1. $x_3x_4^{15}$	2. $x_3^3x_4^{13}$	3. $x_3^{15}x_4$	4. $x_2x_4^{15}$	5. $x_2x_3x_4^{14}$
6. $x_2x_3^2x_4^{13}$	7. $x_2x_3^3x_4^{12}$	8. $x_2x_3^{14}x_4$	9. $x_2x_3^{15}$	10. $x_2^3x_4^{13}$
11. $x_2^3x_3x_4^{12}$	12. $x_2^3x_3^{13}$	13. $x_2^{15}x_4$	14. $x_2^{15}x_3$	15. $x_1x_4^{15}$
16. $x_1x_3x_4^{14}$	17. $x_1x_3^2x_4^{13}$	18. $x_1x_3^3x_4^{12}$	19. $x_1x_3^{14}x_4$	20. $x_1x_3^{15}$
21. $x_1x_2x_4^{14}$	22. $x_1x_2x_3^2x_4^{12}$	23. $x_1x_2x_3^{14}$	24. $x_1x_2^2x_4^{13}$	25. $x_1x_2^2x_3x_4^{12}$
26. $x_1x_2^2x_3^4x_4^9$	27. $x_1x_2^2x_3^5x_4^8$	28. $x_1x_2^2x_3^{12}x_4$	29. $x_1x_2^2x_3^{13}$	30. $x_1x_2^3x_4^{12}$
31. $x_1x_2^3x_3^4x_4^8$	32. $x_1x_2^3x_3^{12}$	33. $x_1x_2^{14}x_4$	34. $x_1x_2^{14}x_3$	35. $x_1x_2^{15}$
36. $x_1^3x_4^{13}$	37. $x_1^3x_3x_4^{12}$	38. $x_1^3x_3^{13}$	39. $x_1^3x_2x_4^{12}$	40. $x_1^3x_2x_3^4x_4^8$
41. $x_1^3x_2x_3^{12}$	42. $x_1^3x_2^{13}$	43. $x_1^{15}x_4$	44. $x_1^{15}x_3$	45. $x_1^{15}x_2$

46. $x_1x_2^3x_3^6x_4^6$ 47. $x_1^3x_2x_3^6x_4^6$ 48. $x_1^3x_2^5x_3^2x_4^6$ 49. $x_1^3x_2^5x_3^6x_4^2$ 50. $x_1x_2x_3^7x_4^7$
51. $x_1x_2^3x_3^5x_4^7$ 52. $x_1x_2^3x_3^7x_4^5$ 53. $x_1x_2^7x_3x_4^7$ 54. $x_1x_2^7x_3^3x_4^5$ 55. $x_1x_2^7x_3^7x_4$
56. $x_1^3x_2x_3^5x_4^7$ 57. $x_1^3x_2x_3^7x_4^5$ 58. $x_1^3x_2^3x_3^5x_4^5$ 59. $x_1^3x_2^5x_3x_4^7$ 60. $x_1^3x_2^5x_3^3x_4^5$
61. $x_1^3x_2^5x_3^7x_4$ 62. $x_1^3x_2^7x_3x_4^5$ 63. $x_1^3x_2^7x_3^5x_4$ 64. $x_1^7x_2x_3x_4^7$ 65. $x_1^7x_2x_3^3x_4^5$
66. $x_1^7x_2x_3^7x_4$ 67. $x_1^7x_2^3x_3x_4^5$ 68. $x_1^7x_2^3x_3^5x_4$ 69. $x_1^7x_2^7x_3x_4$ 70. $x_1^3x_2^3x_3^3x_4^7$
71. $x_1^3x_2^3x_3^7x_4^3$ 72. $x_1^3x_2^7x_3^3x_4^3$ 73. $x_1^7x_2^3x_3^3x_4^3$.

$B_5(16) = B_5^0(16) \cup B_5^+(16)$, where $B_5^0(156) = f(B_4(16))$, $|B_5^0(16)| = 255$ and

$B_5^+(16) = B_5^+(2,1,1,1) \cup B_5^+(2,1,3) \cup B_5^+(2,3,2) \cup B_5^+(2,2,2) \cup B_5^+(4,4,1)$.

6.4.2. $B_5^+(2,1,1,1)$ is the set of 4 monomials $a_t = a_{16,t},\ 1 \leqslant t \leqslant 4$:

1. $x_1x_2x_3^2x_4^4x_5^8$ 2. $x_1x_2^2x_3x_4^4x_5^8$ 3. $x_1x_2^2x_3^4x_4x_5^8$ 4. $x_1x_2^2x_3^4x_4^8x_5$.

6.4.3. $B_5^+(2,1,3)$ is the set of 5 monomials $a_t = a_{16,t},\ 5 \leqslant t \leqslant 9$:

5. $x_1x_2^2x_3^4x_4^4x_5^5$ 6. $x_1x_2^2x_3^4x_4^5x_5^4$ 7. $x_1x_2^2x_3^5x_4^4x_5^4$ 8. $x_1x_2^3x_3^4x_4^4x_5^4$
9. $x_1^3x_2x_3^4x_4^4x_5^4$.

6.4.4. $B_5^+(2,3,2)$ is the set of 20 monomials $a_t = a_{16,t},\ 10 \leqslant t \leqslant 29$:

10. $x_1x_2x_3^2x_4^6x_5^6$ 11. $x_1x_2x_3^6x_4^2x_5^6$ 12. $x_1x_2x_3^6x_4^6x_5^2$ 13. $x_1x_2^2x_3x_4^6x_5^6$
14. $x_1x_2^2x_3^5x_4^2x_5^6$ 15. $x_1x_2^2x_3^5x_4^6x_5^2$ 16. $x_1x_2^3x_3^2x_4^4x_5^6$ 17. $x_1x_2^3x_3^2x_4^6x_5^4$
18. $x_1x_2^3x_3^4x_4^2x_5^6$ 19. $x_1x_2^3x_3^4x_4^6x_5^2$ 20. $x_1x_2^3x_3^6x_4^2x_5^4$ 21. $x_1x_2^3x_3^6x_4^4x_5^2$
22. $x_1^3x_2x_3^2x_4^4x_5^6$ 23. $x_1^3x_2x_3^2x_4^6x_5^4$ 24. $x_1^3x_2x_3^4x_4^2x_5^6$ 25. $x_1^3x_2x_3^4x_4^6x_5^2$
26. $x_1^3x_2x_3^6x_4^2x_5^4$ 27. $x_1^3x_2x_3^6x_4^4x_5^2$ 28. $x_1^3x_2^5x_3^2x_4^2x_5^4$ 29. $x_1^3x_2^5x_3^2x_4^4x_5^2$.

6.4.5. $B_5^+(4,2,2)$ is the set of 110 monomials $a_t = a_{16,t},\ 30 \leqslant t \leqslant 139$:

30. $x_1x_2x_3x_4^6x_5^7$ 31. $x_1x_2x_3x_4^7x_5^6$ 32. $x_1x_2x_3^6x_4x_5^7$ 33. $x_1x_2x_3^6x_4^7x_5$
34. $x_1x_2x_3^7x_4x_5^6$ 35. $x_1x_2x_3^7x_4^6x_5$ 36. $x_1x_2^6x_3x_4x_5^7$ 37. $x_1x_2^6x_3x_4^7x_5$
38. $x_1x_2^6x_3^7x_4x_5$ 39. $x_1x_2^7x_3x_4x_5^6$ 40. $x_1x_2^7x_3x_4^6x_5$ 41. $x_1x_2^7x_3^6x_4x_5$
42. $x_1^7x_2x_3x_4x_5^6$ 43. $x_1^7x_2x_3x_4^6x_5$ 44. $x_1^7x_2x_3^6x_4x_5$ 45. $x_1x_2x_3^2x_4^5x_5^7$
46. $x_1x_2x_3^2x_4^7x_5^5$ 47. $x_1x_2x_3^7x_4^2x_5^5$ 48. $x_1x_2^2x_3x_4^5x_5^7$ 49. $x_1x_2^2x_3x_4^7x_5^5$
50. $x_1x_2^2x_3^5x_4x_5^7$ 51. $x_1x_2^2x_3^5x_4^7x_5$ 52. $x_1x_2^2x_3^7x_4x_5^5$ 53. $x_1x_2^2x_3^7x_4^5x_5$
54. $x_1x_2^7x_3x_4^2x_5^5$ 55. $x_1x_2^7x_3^2x_4x_5^5$ 56. $x_1x_2^7x_3^2x_4^5x_5$ 57. $x_1^7x_2x_3x_4^2x_5^5$
58. $x_1^7x_2x_3^2x_4x_5^5$ 59. $x_1^7x_2x_3^2x_4^5x_5$ 60. $x_1x_2x_3^3x_4^4x_5^7$ 61. $x_1x_2x_3^7x_4^4x_5^4$
62. $x_1x_2x_3^7x_4^3x_5^4$ 63. $x_1x_2^3x_3x_4^4x_5^7$ 64. $x_1x_2^3x_3x_4^7x_5^4$ 65. $x_1x_2^3x_3^4x_4x_5^7$
66. $x_1x_2^3x_3^4x_4^7x_5$ 67. $x_1x_2^3x_3^7x_4x_5^4$ 68. $x_1x_2^3x_3^7x_4^4x_5$ 69. $x_1x_2^7x_3x_4^3x_5^4$
70. $x_1x_2^7x_3^3x_4x_5^4$ 71. $x_1x_2^7x_3^3x_4^4x_5$ 72. $x_1^3x_2x_3x_4^4x_5^7$ 73. $x_1^3x_2x_3x_4^7x_5^4$
74. $x_1^3x_2x_3^4x_4x_5^7$ 75. $x_1^3x_2x_3^4x_4^7x_5$ 76. $x_1^3x_2x_3^7x_4x_5^4$ 77. $x_1^3x_2x_3^7x_4^4x_5$
78. $x_1^3x_2^4x_3x_4x_5^7$ 79. $x_1^3x_2^4x_3x_4^7x_5$ 80. $x_1^3x_2^4x_3^7x_4x_5$ 81. $x_1^3x_2^7x_3x_4x_5^4$
82. $x_1^3x_2^7x_3x_4^4x_5$ 83. $x_1^3x_2^7x_3^4x_4x_5$ 84. $x_1^7x_2x_3x_4^3x_5^4$ 85. $x_1^7x_2x_3^3x_4x_5^4$
86. $x_1^7x_2x_3^3x_4^4x_5$ 87. $x_1^7x_2^3x_3x_4x_5^4$ 88. $x_1^7x_2^3x_3x_4^4x_5$ 89. $x_1^7x_2^3x_3^4x_4x_5$
90. $x_1x_2x_3^3x_4^5x_5^6$ 91. $x_1x_2x_3^3x_4^6x_5^5$ 92. $x_1x_2x_3^6x_4^3x_5^5$ 93. $x_1x_2^3x_3x_4^5x_5^6$
94. $x_1x_2^3x_3x_4^6x_5^5$ 95. $x_1x_2^3x_3^5x_4x_5^6$ 96. $x_1x_2^3x_3^5x_4^6x_5$ 97. $x_1x_2^3x_3^6x_4x_5^5$
98. $x_1x_2^3x_3^6x_4^5x_5$ 99. $x_1x_2^6x_3x_4^3x_5^5$ 100. $x_1x_2^6x_3^3x_4x_5^5$ 101. $x_1x_2^6x_3^3x_4^5x_5$
102. $x_1^3x_2x_3x_4^5x_5^6$ 103. $x_1^3x_2x_3x_4^6x_5^5$ 104. $x_1^3x_2x_3^5x_4x_5^6$ 105. $x_1^3x_2x_3^5x_4^6x_5$
106. $x_1^3x_2x_3^6x_4x_5^5$ 107. $x_1^3x_2x_3^6x_4^5x_5$ 108. $x_1^3x_2^5x_3x_4x_5^6$ 109. $x_1^3x_2^5x_3x_4^6x_5$

110. $x_1^3x_2^5x_3^6x_4x_5$ 111. $x_1x_2^2x_3^3x_4^5x_5^5$ 112. $x_1x_2^2x_3^5x_4^3x_5^5$ 113. $x_1x_2^3x_3^2x_4^5x_5^5$
114. $x_1x_2^3x_3^5x_4^2x_5^5$ 115. $x_1^3x_2x_3^2x_4^5x_5^5$ 116. $x_1^3x_2x_3^5x_4^2x_5^5$ 117. $x_1^3x_2^5x_3x_4^2x_5^5$
118. $x_1^3x_2^5x_3^2x_4x_5^5$ 119. $x_1^3x_2^5x_3^2x_4^5x_5$ 120. $x_1x_2^3x_3^3x_4^4x_5^5$ 121. $x_1x_2^3x_3^3x_4^5x_5^4$
122. $x_1x_2^3x_3^4x_4^3x_5^5$ 123. $x_1x_2^3x_3^5x_4^3x_5^4$ 124. $x_1^3x_2x_3^3x_4^4x_5^5$ 125. $x_1^3x_2x_3^3x_4^5x_5^4$
126. $x_1^3x_2x_3^4x_4^3x_5^5$ 127. $x_1^3x_2x_3^5x_4^3x_5^4$ 128. $x_1^3x_2^3x_3x_4^4x_5^5$ 129. $x_1^3x_2^3x_3x_4^5x_5^4$
130. $x_1^3x_2^3x_3^4x_4x_5^5$ 131. $x_1^3x_2^3x_3^4x_4^5x_5$ 132. $x_1^3x_2^3x_3^5x_4x_5^4$ 133. $x_1^3x_2^3x_3^5x_4^4x_5$
134. $x_1^3x_2^4x_3x_4^3x_5^5$ 135. $x_1^3x_2^4x_3^3x_4x_5^5$ 136. $x_1^3x_2^4x_3^3x_4^5x_5$ 137. $x_1^3x_2^5x_3x_4^3x_5^4$
138. $x_1^3x_2^5x_3^3x_4x_5^4$ 139. $x_1^3x_2^5x_3^3x_4^4x_5$.

6.4.6. $B_5^+(4,4,1)$ is the set of 49 monomials $a_t = a_{16,t},\ 140 \leqslant t \leqslant 188$:

140. $x_1x_2^2x_3^3x_4^3x_5^7$ 141. $x_1x_2^2x_3^3x_4^7x_5^3$ 142. $x_1x_2^2x_3^7x_4^3x_5^3$ 143. $x_1x_2^3x_3^2x_4^3x_5^7$
144. $x_1x_2^3x_3^2x_4^7x_5^3$ 145. $x_1x_2^3x_3^3x_4^2x_5^7$ 146. $x_1x_2^3x_3^3x_4^7x_5^2$ 147. $x_1x_2^3x_3^7x_4^2x_5^3$
148. $x_1x_2^3x_3^7x_4^3x_5^2$ 149. $x_1x_2^7x_3^2x_4^3x_5^3$ 150. $x_1x_2^7x_3^3x_4^2x_5^3$ 151. $x_1x_2^7x_3^3x_4^3x_5^2$
152. $x_1^3x_2x_3^2x_4^3x_5^7$ 153. $x_1^3x_2x_3^2x_4^7x_5^3$ 154. $x_1^3x_2x_3^3x_4^2x_5^7$ 155. $x_1^3x_2x_3^3x_4^7x_5^2$
156. $x_1^3x_2x_3^7x_4^2x_5^3$ 157. $x_1^3x_2x_3^7x_4^3x_5^2$ 158. $x_1^3x_2^3x_3x_4^2x_5^7$ 159. $x_1^3x_2^3x_3x_4^7x_5^2$
160. $x_1^3x_2^3x_3^7x_4x_5^2$ 161. $x_1^3x_2^7x_3x_4^2x_5^3$ 162. $x_1^3x_2^7x_3x_4^3x_5^2$ 163. $x_1^3x_2^7x_3^2x_4x_5^2$
164. $x_1^7x_2x_3^2x_4^3x_5^3$ 165. $x_1^7x_2x_3^3x_4^2x_5^3$ 166. $x_1^7x_2x_3^3x_4^3x_5^2$ 167. $x_1^7x_2^3x_3x_4^2x_5^3$
168. $x_1^7x_2^3x_3x_4^3x_5^2$ 169. $x_1^7x_2^3x_3^3x_4x_5^2$ 170. $x_1x_2^3x_3^3x_4^6x_5^3$ 171. $x_1x_2^3x_3^6x_4^3x_5^3$
172. $x_1x_2^3x_3^6x_4^3x_5^3$ 173. $x_1x_2^6x_3^3x_4^3x_5^3$ 174. $x_1^3x_2x_3^3x_4^3x_5^6$ 175. $x_1^3x_2x_3^3x_4^6x_5^3$
176. $x_1^3x_2x_3^6x_4^3x_5^3$ 177. $x_1^3x_2^3x_3x_4^3x_5^6$ 178. $x_1^3x_2^3x_3x_4^6x_5^3$ 179. $x_1^3x_2^3x_3^3x_4x_5^6$
180. $x_1^3x_2^3x_3^3x_4^5x_5^2$ 181. $x_1^3x_2^3x_3^5x_4^2x_5^3$ 182. $x_1^3x_2^3x_3^5x_4^3x_5^2$ 183. $x_1^3x_2^5x_3^2x_4^3x_5^3$
184. $x_1^3x_2^5x_3^3x_4^2x_5^3$ 185. $x_1^3x_2^5x_3^3x_4^3x_5^2$ 186. $x_1^3x_2^3x_3^3x_4^3x_5^4$ 187. $x_1^3x_2^3x_3^3x_4^4x_5^3$
188. $x_1^3x_2^3x_3^4x_4^3x_5^3$.

6.5. **The admissible monomials of degree 35 in $\mathbf{P_5^0}$.**

6.5.1. $B_5^0(35)$ is the set of 460 monomials, $b_t = b_{35,t},\ 1 \leqslant t \leqslant 460$, determined as follows:

1. $x_3x_4^3x_5^{31}$ 2. $x_3x_4^{31}x_5^3$ 3. $x_3^3x_4x_5^{31}$ 4. $x_3^3x_4^{31}x_5$
5. $x_3^{31}x_4x_5^3$ 6. $x_3^{31}x_4^3x_5$ 7. $x_2x_4^3x_5^{31}$ 8. $x_2x_4^{31}x_5^3$
9. $x_2x_3^3x_5^{31}$ 10. $x_2x_3^3x_4^{31}$ 11. $x_2x_3^{31}x_5^3$ 12. $x_2x_3^{31}x_4^3$
13. $x_2^3x_4x_5^{31}$ 14. $x_2^3x_4^{31}x_5$ 15. $x_2^3x_3x_5^{31}$ 16. $x_2^3x_3x_4^{31}$
17. $x_2^3x_3^{31}x_5$ 18. $x_2^3x_3^{31}x_4$ 19. $x_2^{31}x_4x_5^3$ 20. $x_2^{31}x_4^3x_5$
21. $x_2^{31}x_3x_5^3$ 22. $x_2^{31}x_3x_4^3$ 23. $x_2^{31}x_3^3x_5$ 24. $x_2^{31}x_3^3x_4$
25. $x_1x_4^3x_5^{31}$ 26. $x_1x_4^{31}x_5^3$ 27. $x_1x_3^3x_5^{31}$ 28. $x_1x_3^3x_4^{31}$
29. $x_1x_3^{31}x_5^3$ 30. $x_1x_3^{31}x_4^3$ 31. $x_1x_2^3x_5^{31}$ 32. $x_1x_2^3x_4^{31}$
33. $x_1x_2^3x_3^{31}$ 34. $x_1x_2^{31}x_5^3$ 35. $x_1x_2^{31}x_4^3$ 36. $x_1x_2^{31}x_3^3$
37. $x_1^3x_4x_5^{31}$ 38. $x_1^3x_4^{31}x_5$ 39. $x_1^3x_3x_5^{31}$ 40. $x_1^3x_3x_4^{31}$
41. $x_1^3x_3^{31}x_5$ 42. $x_1^3x_3^{31}x_4$ 43. $x_1^3x_2x_5^{31}$ 44. $x_1^3x_2x_4^{31}$
45. $x_1^3x_2x_3^{31}$ 46. $x_1^3x_2^{31}x_5$ 47. $x_1^3x_2^{31}x_4$ 48. $x_1^3x_2^{31}x_3$
49. $x_1^{31}x_4x_5^3$ 50. $x_1^{31}x_4^3x_5$ 51. $x_1^{31}x_3x_5^3$ 52. $x_1^{31}x_3x_4^3$
53. $x_1^{31}x_3^3x_5$ 54. $x_1^{31}x_3^3x_4$ 55. $x_1^{31}x_2x_5^3$ 56. $x_1^{31}x_2x_4^3$
57. $x_1^{31}x_2x_3^3$ 58. $x_1^{31}x_2^3x_5$ 59. $x_1^{31}x_2^3x_4$ 60. $x_1^{31}x_2^3x_3$
61. $x_3x_4^7x_5^{27}$ 62. $x_3^7x_4x_5^{27}$ 63. $x_3^7x_4^{27}x_5$ 64. $x_2x_4^7x_5^{27}$
65. $x_2x_3^7x_5^{27}$ 66. $x_2x_3^7x_4^{27}$ 67. $x_2^7x_4x_5^{27}$ 68. $x_2^7x_4^{27}x_5$
69. $x_2^7x_3x_5^{27}$ 70. $x_2^7x_3x_4^{27}$ 71. $x_2^7x_3^{27}x_5$ 72. $x_2^7x_3^{27}x_4$
73. $x_1x_4^7x_5^{27}$ 74. $x_1x_3^7x_5^{27}$ 75. $x_1x_3^7x_4^{27}$ 76. $x_1x_2^7x_5^{27}$
77. $x_1x_2^7x_4^{27}$ 78. $x_1x_2^7x_3^{27}$ 79. $x_1^7x_4x_5^{27}$ 80. $x_1^7x_4^{27}x_5$

81. $x_1^7x_3x_5^{27}$ 82. $x_1^7x_3x_4^{27}$ 83. $x_1^7x_3^{27}x_5$ 84. $x_1^7x_3^{27}x_4$
85. $x_1^7x_2x_5^{27}$ 86. $x_1^7x_2x_4^{27}$ 87. $x_1^7x_2x_3^{27}$ 88. $x_1^7x_2^{27}x_5$
89. $x_1^7x_2^{27}x_4$ 90. $x_1^7x_2^{27}x_3$ 91. $x_3^3x_4^3x_5^{29}$ 92. $x_3^3x_4^{29}x_5^3$
93. $x_2^3x_4^3x_5^{29}$ 94. $x_2^3x_4^{29}x_5^3$ 95. $x_2^3x_3^3x_5^{29}$ 96. $x_2^3x_3^3x_4^{29}$
97. $x_2^3x_3^{29}x_5^3$ 98. $x_2^3x_3^{29}x_4^3$ 99. $x_1^3x_4^3x_5^{29}$ 100. $x_1^3x_4^{29}x_5^3$
101. $x_1^3x_3^3x_5^{29}$ 102. $x_1^3x_3^3x_4^{29}$ 103. $x_1^3x_3^{29}x_5^3$ 104. $x_1^3x_3^{29}x_4^3$
105. $x_1^3x_2^3x_5^{29}$ 106. $x_1^3x_2^3x_4^{29}$ 107. $x_1^3x_2^3x_3^{29}$ 108. $x_1^3x_2^{29}x_5^3$
109. $x_1^3x_2^{29}x_4^3$ 110. $x_1^3x_2^{29}x_3^3$ 111. $x_3^3x_4^5x_5^{27}$ 112. $x_2^3x_4^5x_5^{27}$
113. $x_2^3x_3^5x_5^{27}$ 114. $x_2^3x_3^5x_4^{27}$ 115. $x_1^3x_4^5x_5^{27}$ 116. $x_1^3x_3^5x_5^{27}$
117. $x_1^3x_3^5x_4^{27}$ 118. $x_1^3x_2^5x_5^{27}$ 119. $x_1^3x_2^5x_4^{27}$ 120. $x_1^3x_2^5x_3^{27}$
121. $x_3^3x_4^7x_5^{25}$ 122. $x_3^7x_4^3x_5^{25}$ 123. $x_2^3x_4^7x_5^{25}$ 124. $x_2^3x_3^7x_5^{25}$
125. $x_2^3x_3^7x_4^{25}$ 126. $x_2^7x_4^3x_5^{25}$ 127. $x_2^7x_3^3x_5^{25}$ 128. $x_2^7x_3^3x_4^{25}$
129. $x_1^3x_4^7x_5^{25}$ 130. $x_1^3x_3^7x_5^{25}$ 131. $x_1^3x_3^7x_4^{25}$ 132. $x_1^3x_2^7x_5^{25}$
133. $x_1^3x_2^7x_4^{25}$ 134. $x_1^3x_2^7x_3^{25}$ 135. $x_1^7x_4^3x_5^{25}$ 136. $x_1^7x_3^3x_5^{25}$
137. $x_1^7x_3^3x_4^{25}$ 138. $x_1^7x_2^3x_5^{25}$ 139. $x_1^7x_2^3x_4^{25}$ 140. $x_1^7x_2^3x_3^{25}$
141. $x_2x_3x_4^2x_5^{31}$ 142. $x_2x_3x_4^{31}x_5^2$ 143. $x_2x_3^2x_4x_5^{31}$ 144. $x_2x_3^2x_4^{31}x_5$
145. $x_2x_3^{31}x_4x_5^2$ 146. $x_2x_3^{31}x_4^2x_5$ 147. $x_2^{31}x_3x_4x_5^2$ 148. $x_2^{31}x_3x_4^2x_5$
149. $x_1x_3x_4^2x_5^{31}$ 150. $x_1x_3x_4^{31}x_5^2$ 151. $x_1x_3^2x_4x_5^{31}$ 152. $x_1x_3^2x_4^{31}x_5$
153. $x_1x_3^{31}x_4x_5^2$ 154. $x_1x_3^{31}x_4^2x_5$ 155. $x_1x_2x_4^2x_5^{31}$ 156. $x_1x_2x_4^{31}x_5^2$
157. $x_1x_2x_3^2x_5^{31}$ 158. $x_1x_2x_3^2x_4^{31}$ 159. $x_1x_2x_3^{31}x_5^2$ 160. $x_1x_2x_3^{31}x_4^2$
161. $x_1x_2^2x_4x_5^{31}$ 162. $x_1x_2^2x_4^{31}x_5$ 163. $x_1x_2^2x_3x_5^{31}$ 164. $x_1x_2^2x_3x_4^{31}$
165. $x_1x_2^2x_3^{31}x_5$ 166. $x_1x_2^2x_3^{31}x_4$ 167. $x_1x_2^{31}x_4x_5^2$ 168. $x_1x_2^{31}x_4^2x_5$
169. $x_1x_2^{31}x_3x_5^2$ 170. $x_1x_2^{31}x_3x_4^2$ 171. $x_1x_2^{31}x_3^2x_5$ 172. $x_1x_2^{31}x_3^2x_4$
173. $x_1^{31}x_3x_4x_5^2$ 174. $x_1^{31}x_3x_4^2x_5$ 175. $x_1^{31}x_2x_4x_5^2$ 176. $x_1^{31}x_2x_4^2x_5$
177. $x_1^{31}x_2x_3x_5^2$ 178. $x_1^{31}x_2x_3x_4^2$ 179. $x_1^{31}x_2x_3^2x_5$ 180. $x_1^{31}x_2x_3^2x_4$
181. $x_2x_3x_4^3x_5^{30}$ 182. $x_2x_3x_4^6x_5^{27}$ 183. $x_2x_3x_4^7x_5^{26}$ 184. $x_2x_3x_4^{30}x_5^3$
185. $x_2x_3^2x_4^3x_5^{29}$ 186. $x_2x_3^2x_4^5x_5^{27}$ 187. $x_2x_3^2x_4^7x_5^{25}$ 188. $x_2x_3^2x_4^{29}x_5^3$
189. $x_2x_3^3x_4x_5^{30}$ 190. $x_2x_3^3x_4^2x_5^{29}$ 191. $x_2x_3^3x_4^3x_5^{28}$ 192. $x_2x_3^3x_4^4x_5^{27}$
193. $x_2x_3^3x_4^5x_5^{26}$ 194. $x_2x_3^3x_4^6x_5^{25}$ 195. $x_2x_3^3x_4^7x_5^{24}$ 196. $x_2x_3^3x_4^{28}x_5^3$
197. $x_2x_3^3x_4^{29}x_5^2$ 198. $x_2x_3^3x_4^{30}x_5$ 199. $x_2x_3^6x_4x_5^{27}$ 200. $x_2x_3^6x_4^3x_5^{25}$
201. $x_2x_3^6x_4^{27}x_5$ 202. $x_2x_3^7x_4x_5^{26}$ 203. $x_2x_3^7x_4^2x_5^{25}$ 204. $x_2x_3^7x_4^3x_5^{24}$
205. $x_2x_3^7x_4^{26}x_5$ 206. $x_2x_3^{30}x_4x_5^3$ 207. $x_2x_3^{30}x_4^3x_5$ 208. $x_2^3x_3x_4x_5^{30}$
209. $x_2^3x_3x_4^2x_5^{29}$ 210. $x_2^3x_3x_4^3x_5^{28}$ 211. $x_2^3x_3x_4^4x_5^{27}$ 212. $x_2^3x_3x_4^5x_5^{26}$
213. $x_2^3x_3x_4^6x_5^{25}$ 214. $x_2^3x_3x_4^7x_5^{24}$ 215. $x_2^3x_3x_4^{28}x_5^3$ 216. $x_2^3x_3x_4^{29}x_5^2$
217. $x_2^3x_3x_4^{30}x_5$ 218. $x_2^3x_3^3x_4x_5^{28}$ 219. $x_2^3x_3^3x_4^4x_5^{25}$ 220. $x_2^3x_3^3x_4^5x_5^{24}$
221. $x_2^3x_3^3x_4^{28}x_5$ 222. $x_2^3x_3^4x_4x_5^{27}$ 223. $x_2^3x_3^4x_4^3x_5^{25}$ 224. $x_2^3x_3^4x_4^{27}x_5$
225. $x_2^3x_3^5x_4x_5^{26}$ 226. $x_2^3x_3^5x_4^2x_5^{25}$ 227. $x_2^3x_3^5x_4^3x_5^{24}$ 228. $x_2^3x_3^5x_4^{26}x_5$
229. $x_2^3x_3^7x_4x_5^{24}$ 230. $x_2^3x_3^{29}x_4x_5^2$ 231. $x_2^3x_3^{29}x_4^2x_5$ 232. $x_2^7x_3x_4x_5^{26}$
233. $x_2^7x_3x_4^2x_5^{25}$ 234. $x_2^7x_3x_4^3x_5^{24}$ 235. $x_2^7x_3x_4^{26}x_5$ 236. $x_2^7x_3^3x_4x_5^{24}$
237. $x_1x_3x_4^3x_5^{30}$ 238. $x_1x_3x_4^6x_5^{27}$ 239. $x_1x_3x_4^7x_5^{26}$ 240. $x_1x_3x_4^{30}x_5^3$
241. $x_1x_3^2x_4^3x_5^{29}$ 242. $x_1x_3^2x_4^5x_5^{27}$ 243. $x_1x_3^2x_4^7x_5^{25}$ 244. $x_1x_3^2x_4^{29}x_5^3$
245. $x_1x_3^3x_4x_5^{30}$ 246. $x_1x_3^3x_4^2x_5^{29}$ 247. $x_1x_3^3x_4^3x_5^{28}$ 248. $x_1x_3^3x_4^4x_5^{27}$
249. $x_1x_3^3x_4^5x_5^{26}$ 250. $x_1x_3^3x_4^6x_5^{25}$ 251. $x_1x_3^3x_4^7x_5^{24}$ 252. $x_1x_3^3x_4^{28}x_5^3$
253. $x_1x_3^3x_4^{29}x_5^2$ 254. $x_1x_3^3x_4^{30}x_5$ 255. $x_1x_3^6x_4x_5^{27}$ 256. $x_1x_3^6x_4^3x_5^{25}$
257. $x_1x_3^6x_4^{27}x_5$ 258. $x_1x_3^7x_4x_5^{26}$ 259. $x_1x_3^7x_4^2x_5^{25}$ 260. $x_1x_3^7x_4^3x_5^{24}$
261. $x_1x_3^7x_4^{26}x_5$ 262. $x_1x_3^{30}x_4x_5^3$ 263. $x_1x_3^{30}x_4^3x_5$ 264. $x_1x_2x_4^3x_5^{30}$
265. $x_1x_2x_4^6x_5^{27}$ 266. $x_1x_2x_4^7x_5^{26}$ 267. $x_1x_2x_4^{30}x_5^3$ 268. $x_1x_2x_3^3x_5^{30}$
269. $x_1x_2x_3^3x_4^{30}$ 270. $x_1x_2x_3^6x_5^{27}$ 271. $x_1x_2x_3^6x_4^{27}$ 272. $x_1x_2x_3^7x_5^{26}$

273.	$x_1x_2x_3^7x_4^{26}$	274.	$x_1x_2x_3^{30}x_5^3$	275.	$x_1x_2x_3^{30}x_4^3$	276.	$x_1x_2^2x_4^3x_5^{29}$
277.	$x_1x_2^2x_4^5x_5^{27}$	278.	$x_1x_2^2x_4^7x_5^{25}$	279.	$x_1x_2^2x_4^{29}x_5^3$	280.	$x_1x_2^2x_3^3x_5^{29}$
281.	$x_1x_2^2x_3^3x_4^{29}$	282.	$x_1x_2^2x_3^5x_5^{27}$	283.	$x_1x_2^2x_3^5x_4^{27}$	284.	$x_1x_2^2x_3^7x_5^{25}$
285.	$x_1x_2^2x_3^7x_4^{25}$	286.	$x_1x_2^2x_3^{29}x_5^3$	287.	$x_1x_2^2x_3^{29}x_4^3$	288.	$x_1x_2^3x_4x_5^{30}$
289.	$x_1x_2^3x_4^2x_5^{29}$	290.	$x_1x_2^3x_4^3x_5^{28}$	291.	$x_1x_2^3x_4^4x_5^{27}$	292.	$x_1x_2^3x_4^5x_5^{26}$
293.	$x_1x_2^3x_4^6x_5^{25}$	294.	$x_1x_2^3x_4^7x_5^{24}$	295.	$x_1x_2^3x_4^{28}x_5^3$	296.	$x_1x_2^3x_4^{29}x_5^2$
297.	$x_1x_2^3x_4^{30}x_5$	298.	$x_1x_2^3x_3x_5^{30}$	299.	$x_1x_2^3x_3x_4^{30}$	300.	$x_1x_2^3x_3^2x_5^{29}$
301.	$x_1x_2^3x_3^2x_4^{29}$	302.	$x_1x_2^3x_3^3x_5^{28}$	303.	$x_1x_2^3x_3^3x_4^{28}$	304.	$x_1x_2^3x_3^4x_5^{27}$
305.	$x_1x_2^3x_3^4x_4^{27}$	306.	$x_1x_2^3x_3^5x_5^{26}$	307.	$x_1x_2^3x_3^5x_4^{26}$	308.	$x_1x_2^3x_3^6x_5^{25}$
309.	$x_1x_2^3x_3^6x_4^{25}$	310.	$x_1x_2^3x_3^7x_5^{24}$	311.	$x_1x_2^3x_3^7x_4^{24}$	312.	$x_1x_2^3x_3^{28}x_5^3$
313.	$x_1x_2^3x_3^{28}x_4^3$	314.	$x_1x_2^3x_3^{29}x_5^2$	315.	$x_1x_2^3x_3^{29}x_4^2$	316.	$x_1x_2^3x_3^{30}x_5$
317.	$x_1x_2^3x_3^{30}x_4$	318.	$x_1x_2^6x_4x_5^{27}$	319.	$x_1x_2^6x_4^3x_5^{25}$	320.	$x_1x_2^6x_4^{27}x_5$
321.	$x_1x_2^6x_3x_5^{27}$	322.	$x_1x_2^6x_3x_4^{27}$	323.	$x_1x_2^6x_3^3x_5^{25}$	324.	$x_1x_2^6x_3^3x_4^{25}$
325.	$x_1x_2^6x_3^{27}x_5$	326.	$x_1x_2^6x_3^{27}x_4$	327.	$x_1x_2^7x_4x_5^{26}$	328.	$x_1x_2^7x_4^2x_5^{25}$
329.	$x_1x_2^7x_4^3x_5^{24}$	330.	$x_1x_2^7x_4^{26}x_5$	331.	$x_1x_2^7x_3x_5^{26}$	332.	$x_1x_2^7x_3x_4^{26}$
333.	$x_1x_2^7x_3^2x_5^{25}$	334.	$x_1x_2^7x_3^2x_4^{25}$	335.	$x_1x_2^7x_3^3x_5^{24}$	336.	$x_1x_2^7x_3^3x_4^{24}$
337.	$x_1x_2^7x_3^{26}x_5$	338.	$x_1x_2^7x_3^{26}x_4$	339.	$x_1x_2^{30}x_4x_5^3$	340.	$x_1x_2^{30}x_4^3x_5$
341.	$x_1x_2^{30}x_3x_5^3$	342.	$x_1x_2^{30}x_3x_4^3$	343.	$x_1x_2^{30}x_3^3x_5$	344.	$x_1x_2^{30}x_3^3x_4$
345.	$x_1^3x_3x_4x_5^{30}$	346.	$x_1^3x_3x_4^2x_5^{29}$	347.	$x_1^3x_3x_4^3x_5^{28}$	348.	$x_1^3x_3x_4^4x_5^{27}$
349.	$x_1^3x_3x_4^5x_5^{26}$	350.	$x_1^3x_3x_4^6x_5^{25}$	351.	$x_1^3x_3x_4^7x_5^{24}$	352.	$x_1^3x_3x_4^{28}x_5^3$
353.	$x_1^3x_3x_4^{29}x_5^2$	354.	$x_1^3x_3x_4^{30}x_5$	355.	$x_1^3x_3^3x_4x_5^{28}$	356.	$x_1^3x_3^3x_4^4x_5^{25}$
357.	$x_1^3x_3^3x_4^5x_5^{24}$	358.	$x_1^3x_3^3x_4^{28}x_5$	359.	$x_1^3x_3^4x_4x_5^{27}$	360.	$x_1^3x_3^4x_4^3x_5^{25}$
361.	$x_1^3x_3^4x_4^{27}x_5$	362.	$x_1^3x_3^5x_4x_5^{26}$	363.	$x_1^3x_3^5x_4^2x_5^{25}$	364.	$x_1^3x_3^5x_4^3x_5^{24}$
365.	$x_1^3x_3^5x_4^{26}x_5$	366.	$x_1^3x_3^7x_4x_5^{24}$	367.	$x_1^3x_3^{29}x_4x_5^2$	368.	$x_1^3x_3^{29}x_4^2x_5$
369.	$x_1^3x_2x_4x_5^{30}$	370.	$x_1^3x_2x_4^2x_5^{29}$	371.	$x_1^3x_2x_4^3x_5^{28}$	372.	$x_1^3x_2x_4^4x_5^{27}$
373.	$x_1^3x_2x_4^5x_5^{26}$	374.	$x_1^3x_2x_4^6x_5^{25}$	375.	$x_1^3x_2x_4^7x_5^{24}$	376.	$x_1^3x_2x_4^{28}x_5^3$
377.	$x_1^3x_2x_4^{29}x_5^2$	378.	$x_1^3x_2x_4^{30}x_5$	379.	$x_1^3x_2x_3x_5^{30}$	380.	$x_1^3x_2x_3x_4^{30}$
381.	$x_1^3x_2x_3^2x_5^{29}$	382.	$x_1^3x_2x_3^2x_4^{29}$	383.	$x_1^3x_2x_3^3x_5^{28}$	384.	$x_1^3x_2x_3^3x_4^{28}$
385.	$x_1^3x_2x_3^4x_5^{27}$	386.	$x_1^3x_2x_3^4x_4^{27}$	387.	$x_1^3x_2x_3^5x_5^{26}$	388.	$x_1^3x_2x_3^5x_4^{26}$
389.	$x_1^3x_2x_3^6x_5^{25}$	390.	$x_1^3x_2x_3^6x_4^{25}$	391.	$x_1^3x_2x_3^7x_5^{24}$	392.	$x_1^3x_2x_3^7x_4^{24}$
393.	$x_1^3x_2x_3^{28}x_5^3$	394.	$x_1^3x_2x_3^{28}x_4^3$	395.	$x_1^3x_2x_3^{29}x_5^2$	396.	$x_1^3x_2x_3^{29}x_4^2$
397.	$x_1^3x_2x_3^{30}x_5$	398.	$x_1^3x_2x_3^{30}x_4$	399.	$x_1^3x_2^3x_4x_5^{28}$	400.	$x_1^3x_2^3x_4^4x_5^{25}$
401.	$x_1^3x_2^3x_4^5x_5^{24}$	402.	$x_1^3x_2^3x_4^{28}x_5$	403.	$x_1^3x_2^3x_3x_5^{28}$	404.	$x_1^3x_2^3x_3x_4^{28}$
405.	$x_1^3x_2^3x_3^4x_5^{25}$	406.	$x_1^3x_2^3x_3^4x_4^{25}$	407.	$x_1^3x_2^3x_3^5x_5^{24}$	408.	$x_1^3x_2^3x_3^5x_4^{24}$
409.	$x_1^3x_2^3x_3^{28}x_5$	410.	$x_1^3x_2^3x_3^{28}x_4$	411.	$x_1^3x_2^4x_4x_5^{27}$	412.	$x_1^3x_2^4x_4^3x_5^{25}$
413.	$x_1^3x_2^4x_4^{27}x_5$	414.	$x_1^3x_2^4x_3x_5^{27}$	415.	$x_1^3x_2^4x_3x_4^{27}$	416.	$x_1^3x_2^4x_3^3x_5^{25}$
417.	$x_1^3x_2^4x_3^3x_4^{25}$	418.	$x_1^3x_2^4x_3^{27}x_5$	419.	$x_1^3x_2^4x_3^{27}x_4$	420.	$x_1^3x_2^5x_4x_5^{26}$
421.	$x_1^3x_2^5x_4^2x_5^{25}$	422.	$x_1^3x_2^5x_4^3x_5^{24}$	423.	$x_1^3x_2^5x_4^{26}x_5$	424.	$x_1^3x_2^5x_3x_5^{26}$
425.	$x_1^3x_2^5x_3x_4^{26}$	426.	$x_1^3x_2^5x_3^2x_5^{25}$	427.	$x_1^3x_2^5x_3^2x_4^{25}$	428.	$x_1^3x_2^5x_3^3x_5^{24}$
429.	$x_1^3x_2^5x_3^3x_4^{24}$	430.	$x_1^3x_2^5x_3^{26}x_5$	431.	$x_1^3x_2^5x_3^{26}x_4$	432.	$x_1^3x_2^7x_4x_5^{24}$
433.	$x_1^3x_2^7x_3x_5^{24}$	434.	$x_1^3x_2^7x_3x_4^{24}$	435.	$x_1^3x_2^{29}x_4x_5^2$	436.	$x_1^3x_2^{29}x_4^2x_5$
437.	$x_1^3x_2^{29}x_3x_5^2$	438.	$x_1^3x_2^{29}x_3x_4^2$	439.	$x_1^3x_2^{29}x_3^2x_5$	440.	$x_1^3x_2^{29}x_3^2x_4$
441.	$x_1^7x_3x_4x_5^{26}$	442.	$x_1^7x_3x_4^2x_5^{25}$	443.	$x_1^7x_3x_4^3x_5^{24}$	444.	$x_1^7x_3x_4^{26}x_5$
445.	$x_1^7x_3^3x_4x_5^{24}$	446.	$x_1^7x_2x_4x_5^{26}$	447.	$x_1^7x_2x_4^2x_5^{25}$	448.	$x_1^7x_2x_4^3x_5^{24}$
449.	$x_1^7x_2x_4^{26}x_5$	450.	$x_1^7x_2x_3x_5^{26}$	451.	$x_1^7x_2x_3x_4^{26}$	452.	$x_1^7x_2x_3^2x_5^{25}$
453.	$x_1^7x_2x_3^2x_4^{25}$	454.	$x_1^7x_2x_3^3x_5^{24}$	455.	$x_1^7x_2x_3^3x_4^{24}$	456.	$x_1^7x_2x_3^{26}x_5$
457.	$x_1^7x_2x_3^{26}x_4$	458.	$x_1^7x_2^3x_4x_5^{24}$	459.	$x_1^7x_2^3x_3x_5^{24}$	460.	$x_1^7x_2^3x_3x_4^{24}$

We have $B_5(35) = B_5^0(35) \cup B_5^+(\omega_{(1)}) \cup B_5(\omega_{(3)}) \cup B_5(\omega_{(4)}) \cup B_5(\omega_{(3)}) \cup \psi(B_5(15))$.

6.5.2. $B_5^+(\omega_{(1)})$ is the set of 160 monomials $a_t = a_{35,t},\ 1 \leqslant t \leqslant 160$:

1. $x_1x_2x_3x_4^2x_5^{30}$
2. $x_1x_2x_3x_4^6x_5^{26}$
3. $x_1x_2x_3x_4^{30}x_5^2$
4. $x_1x_2x_3^2x_4x_5^{30}$
5. $x_1x_2x_3^2x_4^2x_5^{29}$
6. $x_1x_2x_3^2x_4^3x_5^{28}$
7. $x_1x_2x_3^2x_4^4x_5^{27}$
8. $x_1x_2x_3^2x_4^5x_5^{26}$
9. $x_1x_2x_3^2x_4^6x_5^{25}$
10. $x_1x_2x_3^2x_4^7x_5^{24}$
11. $x_1x_2x_3^2x_4^{28}x_5^3$
12. $x_1x_2x_3^2x_4^{29}x_5^2$
13. $x_1x_2x_3^2x_4^{30}x_5$
14. $x_1x_2x_3^3x_4^2x_5^{28}$
15. $x_1x_2x_3^3x_4^4x_5^{26}$
16. $x_1x_2x_3^3x_4^6x_5^{24}$
17. $x_1x_2x_3^3x_4^{28}x_5^2$
18. $x_1x_2x_3^6x_4x_5^{26}$
19. $x_1x_2x_3^6x_4^2x_5^{25}$
20. $x_1x_2x_3^6x_4^3x_5^{24}$
21. $x_1x_2x_3^6x_4^{26}x_5$
22. $x_1x_2x_3^7x_4^2x_5^{24}$
23. $x_1x_2x_3^{30}x_4x_5^2$
24. $x_1x_2x_3^{30}x_4^2x_5$
25. $x_1x_2^2x_3x_4x_5^{30}$
26. $x_1x_2^2x_3x_4^2x_5^{29}$
27. $x_1x_2^2x_3x_4^3x_5^{28}$
28. $x_1x_2^2x_3x_4^4x_5^{27}$
29. $x_1x_2^2x_3x_4^5x_5^{26}$
30. $x_1x_2^2x_3x_4^6x_5^{25}$
31. $x_1x_2^2x_3x_4^7x_5^{24}$
32. $x_1x_2^2x_3x_4^{28}x_5^3$
33. $x_1x_2^2x_3x_4^{29}x_5^2$
34. $x_1x_2^2x_3x_4^{30}x_5$
35. $x_1x_2^2x_3^3x_4x_5^{28}$
36. $x_1x_2^2x_3^3x_4^4x_5^{25}$
37. $x_1x_2^2x_3^3x_4^5x_5^{24}$
38. $x_1x_2^2x_3^3x_4^{28}x_5$
39. $x_1x_2^2x_3^4x_4x_5^{27}$
40. $x_1x_2^2x_3^4x_4^3x_5^{25}$
41. $x_1x_2^2x_3^4x_4^9x_5^{19}$
42. $x_1x_2^2x_3^4x_4^{11}x_5^{17}$
43. $x_1x_2^2x_3^4x_4^{25}x_5^3$
44. $x_1x_2^2x_3^4x_4^{27}x_5$
45. $x_1x_2^2x_3^5x_4x_5^{26}$
46. $x_1x_2^2x_3^5x_4^2x_5^{25}$
47. $x_1x_2^2x_3^5x_4^3x_5^{24}$
48. $x_1x_2^2x_3^5x_4^8x_5^{19}$
49. $x_1x_2^2x_3^5x_4^9x_5^{18}$
50. $x_1x_2^2x_3^5x_4^{10}x_5^{17}$
51. $x_1x_2^2x_3^5x_4^{11}x_5^{16}$
52. $x_1x_2^2x_3^5x_4^{24}x_5^3$
53. $x_1x_2^2x_3^5x_4^{25}x_5^2$
54. $x_1x_2^2x_3^5x_4^{26}x_5$
55. $x_1x_2^2x_3^7x_4x_5^{24}$
56. $x_1x_2^2x_3^7x_4^8x_5^{17}$
57. $x_1x_2^2x_3^7x_4^9x_5^{16}$
58. $x_1x_2^2x_3^7x_4^{24}x_5$
59. $x_1x_2^2x_3^{28}x_4x_5^3$
60. $x_1x_2^2x_3^{28}x_4^3x_5$
61. $x_1x_2^2x_3^{29}x_4x_5^2$
62. $x_1x_2^2x_3^{29}x_4^2x_5$
63. $x_1x_2^3x_3x_4^2x_5^{28}$
64. $x_1x_2^3x_3x_4^4x_5^{26}$
65. $x_1x_2^3x_3x_4^6x_5^{24}$
66. $x_1x_2^3x_3x_4^{28}x_5^2$
67. $x_1x_2^3x_3^2x_4x_5^{28}$
68. $x_1x_2^3x_3^2x_4^4x_5^{25}$
69. $x_1x_2^3x_3^2x_4^5x_5^{24}$
70. $x_1x_2^3x_3^2x_4^{28}x_5$
71. $x_1x_2^3x_3^3x_4^4x_5^{24}$
72. $x_1x_2^3x_3^4x_4x_5^{26}$
73. $x_1x_2^3x_3^4x_4^2x_5^{25}$
74. $x_1x_2^3x_3^4x_4^3x_5^{24}$
75. $x_1x_2^3x_3^4x_4^8x_5^{19}$
76. $x_1x_2^3x_3^4x_4^9x_5^{18}$
77. $x_1x_2^3x_3^4x_4^{10}x_5^{17}$
78. $x_1x_2^3x_3^4x_4^{11}x_5^{16}$
79. $x_1x_2^3x_3^4x_4^{24}x_5^3$
80. $x_1x_2^3x_3^4x_4^{25}x_5^2$
81. $x_1x_2^3x_3^4x_4^{26}x_5$
82. $x_1x_2^3x_3^5x_4^2x_5^{24}$
83. $x_1x_2^3x_3^5x_4^8x_5^{18}$
84. $x_1x_2^3x_3^5x_4^{10}x_5^{16}$
85. $x_1x_2^3x_3^5x_4^{24}x_5^2$
86. $x_1x_2^3x_3^6x_4x_5^{24}$
87. $x_1x_2^3x_3^6x_4^8x_5^{17}$
88. $x_1x_2^3x_3^6x_4^9x_5^{16}$
89. $x_1x_2^3x_3^6x_4^{24}x_5$
90. $x_1x_2^3x_3^7x_4^8x_5^{16}$
91. $x_1x_2^3x_3^{28}x_4x_5^2$
92. $x_1x_2^3x_3^{28}x_4^2x_5$
93. $x_1x_2^6x_3x_4x_5^{26}$
94. $x_1x_2^6x_3x_4^2x_5^{25}$
95. $x_1x_2^6x_3x_4^3x_5^{24}$
96. $x_1x_2^6x_3x_4^{26}x_5$
97. $x_1x_2^6x_3^3x_4x_5^{24}$
98. $x_1x_2^7x_3x_4^2x_5^{24}$
99. $x_1x_2^7x_3^2x_4x_5^{24}$
100. $x_1x_2^7x_3^2x_4^8x_5^{17}$
101. $x_1x_2^7x_3^2x_4^9x_5^{16}$
102. $x_1x_2^7x_3^2x_4^{24}x_5$
103. $x_1x_2^7x_3^3x_4^8x_5^{16}$
104. $x_1x_2^{30}x_3x_4x_5^2$
105. $x_1x_2^{30}x_3x_4^2x_5$
106. $x_1^3x_2x_3x_4^2x_5^{28}$
107. $x_1^3x_2x_3x_4^4x_5^{26}$
108. $x_1^3x_2x_3x_4^6x_5^{24}$
109. $x_1^3x_2x_3x_4^{28}x_5^2$
110. $x_1^3x_2x_3^2x_4x_5^{28}$
111. $x_1^3x_2x_3^2x_4^4x_5^{25}$
112. $x_1^3x_2x_3^2x_4^5x_5^{24}$
113. $x_1^3x_2x_3^2x_4^{28}x_5$
114. $x_1^3x_2x_3^3x_4^4x_5^{24}$
115. $x_1^3x_2x_3^4x_4x_5^{26}$
116. $x_1^3x_2x_3^4x_4^2x_5^{25}$
117. $x_1^3x_2x_3^4x_4^3x_5^{24}$
118. $x_1^3x_2x_3^4x_4^8x_5^{19}$
119. $x_1^3x_2x_3^4x_4^9x_5^{18}$
120. $x_1^3x_2x_3^4x_4^{10}x_5^{17}$
121. $x_1^3x_2x_3^4x_4^{11}x_5^{16}$
122. $x_1^3x_2x_3^4x_4^{24}x_5^3$
123. $x_1^3x_2x_3^4x_4^{25}x_5^2$
124. $x_1^3x_2x_3^4x_4^{26}x_5$
125. $x_1^3x_2x_3^5x_4^2x_5^{24}$
126. $x_1^3x_2x_3^5x_4^8x_5^{18}$
127. $x_1^3x_2x_3^5x_4^{10}x_5^{16}$
128. $x_1^3x_2x_3^5x_4^{24}x_5^2$
129. $x_1^3x_2x_3^6x_4x_5^{24}$
130. $x_1^3x_2x_3^6x_4^8x_5^{17}$
131. $x_1^3x_2x_3^6x_4^9x_5^{16}$
132. $x_1^3x_2x_3^6x_4^{24}x_5$
133. $x_1^3x_2x_3^7x_4^8x_5^{16}$
134. $x_1^3x_2x_3^{28}x_4x_5^2$
135. $x_1^3x_2x_3^{28}x_4^2x_5$
136. $x_1^3x_2^3x_3x_4^4x_5^{24}$
137. $x_1^3x_2^3x_3^4x_4x_5^{24}$
138. $x_1^3x_2^3x_3^4x_4^8x_5^{17}$
139. $x_1^3x_2^3x_3^4x_4^9x_5^{16}$
140. $x_1^3x_2^3x_3^4x_4^{24}x_5$
141. $x_1^3x_2^3x_3^5x_4^8x_5^{16}$

142. $x_1^3x_2^4x_3x_4x_5^{26}$	143. $x_1^3x_2^4x_3x_4^2x_5^{25}$	144. $x_1^3x_2^4x_3x_4^3x_5^{24}$
145. $x_1^3x_2^4x_3x_4^{26}x_5$	146. $x_1^3x_2^4x_3^3x_4x_5^{24}$	147. $x_1^3x_2^5x_3x_4^2x_5^{24}$
148. $x_1^3x_2^5x_3^2x_4x_5^{24}$	149. $x_1^3x_2^5x_3^2x_4^8x_5^{17}$	150. $x_1^3x_2^5x_3^2x_4^9x_5^{16}$
151. $x_1^3x_2^5x_3^2x_4^{24}x_5$	152. $x_1^3x_2^5x_3^3x_4^8x_5^{16}$	153. $x_1^3x_2^7x_3x_4^8x_5^{16}$
154. $x_1^7x_2x_3x_4^2x_5^{24}$	155. $x_1^7x_2x_3^2x_4x_5^{24}$	156. $x_1^7x_2x_3^2x_4^8x_5^{17}$
157. $x_1^7x_2x_3^2x_4^9x_5^{16}$	158. $x_1^7x_2x_3^2x_4^{24}x_5$	159. $x_1^7x_2x_3^3x_4^8x_5^{16}$
160. $x_1^7x_2^3x_3x_4^8x_5^{16}$		

6.5.3. $B_5^+(\omega_{(4)})$ is the set of 50 monomials

$$a_t = a_{35,t},\ 161 \leqslant t \leqslant 210$$

161. $x_1x_2^2x_3^7x_4^{11}x_5^{14}$	162. $x_1x_2^7x_3^2x_4^{11}x_5^{14}$	163. $x_1x_2^7x_3^{11}x_4^2x_5^{14}$
164. $x_1x_2^7x_3^{11}x_4^{14}x_5^2$	165. $x_1^7x_2x_3^2x_4^{11}x_5^{14}$	166. $x_1^7x_2x_3^{11}x_4^2x_5^{14}$
167. $x_1^7x_2x_3^{11}x_4^{14}x_5^2$	168. $x_1^7x_2^{11}x_3x_4^2x_5^{14}$	169. $x_1^7x_2^{11}x_3x_4^{14}x_5^2$
170. $x_1^7x_2^{11}x_3^{13}x_4^2x_5^2$	171. $x_1x_2^3x_3^6x_4^{11}x_5^{14}$	172. $x_1^3x_2x_3^6x_4^{11}x_5^{14}$
173. $x_1x_2^3x_3^7x_4^{10}x_5^{14}$	174. $x_1x_2^3x_3^7x_4^{14}x_5^{10}$	175. $x_1x_2^7x_3^3x_4^{10}x_5^{14}$
176. $x_1x_2^7x_3^3x_4^{14}x_5^{10}$	177. $x_1^3x_2x_3^7x_4^{10}x_5^{14}$	178. $x_1^3x_2x_3^7x_4^{14}x_5^{10}$
179. $x_1^3x_2^7x_3x_4^{10}x_5^{14}$	180. $x_1^3x_2^7x_3x_4^{14}x_5^{10}$	181. $x_1^7x_2x_3^3x_4^{10}x_5^{14}$
182. $x_1^7x_2x_3^3x_4^{14}x_5^{10}$	183. $x_1^7x_2^3x_3x_4^{10}x_5^{14}$	184. $x_1^7x_2^3x_3x_4^{14}x_5^{10}$
185. $x_1x_2^7x_3^{11}x_4^6x_5^{10}$	186. $x_1^7x_2x_3^{11}x_4^6x_5^{10}$	187. $x_1^7x_2^{11}x_3x_4^6x_5^{10}$
188. $x_1^3x_2^5x_3^2x_4^{11}x_5^{14}$	189. $x_1^3x_2^5x_3^{11}x_4^2x_5^{14}$	190. $x_1^3x_2^5x_3^{11}x_4^{14}x_5^2$
191. $x_1^3x_2^7x_3^9x_4^2x_5^{14}$	192. $x_1^3x_2^7x_3^9x_4^{14}x_5^2$	193. $x_1^7x_2^3x_3^9x_4^2x_5^{14}$
194. $x_1^7x_2^3x_3^9x_4^{14}x_5^2$	195. $x_1^3x_2^7x_3^{13}x_4^2x_5^{10}$	196. $x_1^3x_2^7x_3^{13}x_4^{10}x_5^2$
197. $x_1^7x_2^3x_3^{13}x_4^2x_5^{10}$	198. $x_1^7x_2^3x_3^{13}x_4^{10}x_5^2$	199. $x_1^7x_2^{11}x_3^5x_4^2x_5^{10}$
200. $x_1^7x_2^{11}x_3^5x_4^{10}x_5^2$	201. $x_1^3x_2^3x_3^5x_4^{10}x_5^{14}$	202. $x_1^3x_2^3x_3^5x_4^{14}x_5^{10}$
203. $x_1^3x_2^5x_3^3x_4^{10}x_5^{14}$	204. $x_1^3x_2^5x_3^3x_4^{14}x_5^{10}$	205. $x_1^3x_2^3x_3^{13}x_4^6x_5^{10}$
206. $x_1^3x_2^5x_3^{11}x_4^6x_5^{10}$	207. $x_1^3x_2^7x_3^5x_4^{10}x_5^{10}$	208. $x_1^7x_2^3x_3^5x_4^{10}x_5^{10}$
209. $x_1^3x_2^7x_3^9x_4^6x_5^{10}$	210. $x_1^7x_2^3x_3^9x_4^6x_5^{10}$	

6.5.4. $B_5^+(\omega_{(5)})$ is the set of 15 monomials

$$a_t = a_{35,t},\ 211 \leqslant t \leqslant 225$$

211. $x_1^3x_2^5x_3^6x_4^6x_5^{15}$	212. $x_1^3x_2^5x_3^6x_4^{15}x_5^6$	213. $x_1^3x_2^5x_3^{15}x_4^6x_5^6$
214. $x_1^3x_2^{15}x_3^5x_4^6x_5^6$	215. $x_1^{15}x_2^3x_3^5x_4^6x_5^6$	216. $x_1^3x_2^5x_3^6x_4^7x_5^{14}$
217. $x_1^3x_2^5x_3^7x_4^6x_5^{14}$	218. $x_1^3x_2^5x_3^7x_4^{14}x_5^6$	219. $x_1^3x_2^7x_3^5x_4^6x_5^{14}$
220. $x_1^3x_2^7x_3^5x_4^{14}x_5^6$	221. $x_1^3x_2^7x_3^{13}x_4^6x_5^6$	222. $x_1^7x_2^3x_3^5x_4^6x_5^{14}$
223. $x_1^7x_2^3x_3^5x_4^{14}x_5^6$	224. $x_1^7x_2^3x_3^{13}x_4^6x_5^6$	225. $x_1^7x_2^{11}x_3^5x_4^6x_5^6$

We have $f(B_4(35)) = QP_5^0(\omega_{(1)})$ and $|f(B_4(35))| = 460$. Hence,

$$\dim QP_5(\omega_{(1)}) = 620.$$

6.6. **The Σ_5-invariants of $QP_5(\omega_{(1)})$.**

6.6.1. *The Σ_5-invariants of $QP_5^0(\omega_{(1)}) = (QP_5^0)_{35}$.*

$(QP_5^0)_{35}^{\Sigma_5} = \langle [p_i] : 1 \leqslant i \leqslant 6\rangle$, where $p_1 = p(u_1) = \sum_{t=1}^{60} b_t$, $p_2 = p(u_2) = \sum_{t=61}^{90} b_t$, with $u_1 = x_1x_2^3x_3^{31}$, $u_2 = x_1x_2^7x_3^{27}$,

$$\begin{aligned}
p_3 = {} & x_3^3x_4^{29}x_5^3 + x_2^3x_4^{29}x_5^3 + x_2^3x_3^{29}x_5^3 + x_2^3x_3^{29}x_4^3 + x_1^3x_4^{29}x_5^3 + x_1^3x_3^{29}x_5^3\\
& + x_1^3x_3^{29}x_4^3 + x_1^3x_2^{29}x_5^3 + x_1^3x_2^{29}x_4^3 + x_1^3x_2^{29}x_3^3 + x_3^3x_4^5x_5^{27} + x_2^3x_4^5x_5^{27}\\
& + x_2^3x_3^5x_5^{27} + x_2^3x_3^5x_4^{27} + x_1^3x_4^5x_5^{27} + x_1^3x_3^5x_5^{27} + x_1^3x_3^5x_4^{27} + x_1^3x_2^5x_5^{27}\\
& + x_1^3x_2^5x_4^{27} + x_1^3x_2^5x_3^{27} + x_3^3x_4^7x_5^{25} + x_3^7x_4^3x_5^{25} + x_2^3x_4^7x_5^{25} + x_2^3x_3^7x_5^{25}\\
& + x_2^3x_3^7x_4^{25} + x_2^7x_4^3x_5^{25} + x_2^7x_3^3x_5^{25} + x_2^7x_3^3x_4^{25} + x_1^3x_4^7x_5^{25} + x_1^3x_3^7x_5^{25}\\
& + x_1^3x_3^7x_4^{25} + x_1^3x_2^7x_5^{25} + x_1^3x_2^7x_4^{25} + x_1^3x_2^7x_3^{25} + x_1^7x_4^3x_5^{25} + x_1^7x_3^3x_5^{25}\\
& + x_1^7x_3^3x_4^{25} + x_1^7x_2^3x_5^{25} + x_1^7x_2^3x_4^{25} + x_1^7x_2^3x_3^{25},
\end{aligned}$$

$$\begin{aligned}
p_4 = {} & x_2x_3x_4^3x_5^{30} + x_1x_3x_4^3x_5^{30} + x_1x_2x_4^3x_5^{30} + x_1x_2x_3^3x_5^{30} + x_1x_2x_3^3x_4^{30}\\
& + x_2^3x_3^{29}x_4x_5^2 + x_2^3x_3^{29}x_4^2x_5 + x_1^3x_3^{29}x_4x_5^2 + x_1^3x_3^{29}x_4^2x_5 + x_1^3x_2^{29}x_4x_5^2\\
& + x_1^3x_2^{29}x_4^2x_5 + x_1^3x_2^{29}x_3x_5^2 + x_1^3x_2^{29}x_3x_4^2 + x_1^3x_2^{29}x_3^2x_5 + x_1^3x_2^{29}x_3^2x_4\\
& + x_2^3x_3x_4^3x_5^{28} + x_1^3x_3x_4^3x_5^{28} + x_1^3x_2x_4^3x_5^{28} + x_1^3x_2x_3^3x_5^{28} + x_1^3x_2x_3^3x_4^{28}\\
& + x_2x_3^3x_4^5x_5^{26} + x_2^3x_3x_4^5x_5^{26} + x_2^3x_3^5x_4x_5^{26} + x_2^3x_3^5x_4^{26}x_5 + x_1x_3^3x_4^5x_5^{26}\\
& + x_1x_2^3x_4^5x_5^{26} + x_1x_2^3x_3^5x_5^{26} + x_1x_2^3x_3^5x_4^{26} + x_1^3x_3x_4^5x_5^{26} + x_1^3x_3^5x_4x_5^{26}\\
& + x_1^3x_3^5x_4^{26}x_5 + x_1^3x_2x_4^5x_5^{26} + x_1^3x_2x_3^5x_5^{26} + x_1^3x_2x_3^5x_4^{26} + x_1^3x_2^5x_4x_5^{26}\\
& + x_1^3x_2^5x_4^{26}x_5 + x_1^3x_2^5x_3x_5^{26} + x_1^3x_2^5x_3x_4^{26} + x_1^3x_2^5x_3^{26}x_5 + x_1^3x_2^5x_3^{26}x_4\\
& + x_2x_3^3x_4^6x_5^{25} + x_2x_3^6x_4^3x_5^{25} + x_1x_3^3x_4^6x_5^{25} + x_1x_3^6x_4^3x_5^{25} + x_1x_3^3x_4^6x_5^{25}\\
& + x_1x_2^3x_3^6x_5^{25} + x_1x_2^3x_3^6x_4^{25} + x_1x_2^6x_4^3x_5^{25} + x_1x_2^6x_3^3x_5^{25} + x_1x_2^6x_3^3x_4^{25}\\
& + x_2^3x_3^3x_4^4x_5^{25} + x_2^3x_3^4x_4^3x_5^{25} + x_1^3x_3^3x_4^4x_5^{25} + x_1^3x_3^4x_4^3x_5^{25} + x_1^3x_2^3x_4^4x_5^{25}\\
& + x_1^3x_2^3x_3^4x_5^{25} + x_1^3x_2^3x_3^4x_4^{25} + x_1^3x_2^4x_4^3x_5^{25} + x_1^3x_2^4x_3^3x_5^{25} + x_1^3x_2^4x_3^3x_4^{25}\\
& + x_2^3x_3^3x_4^5x_5^{24} + x_1^3x_3^3x_4^5x_5^{24} + x_1^3x_2^3x_4^5x_5^{24} + x_1^3x_2^3x_3^5x_5^{24} + x_1^3x_2^3x_3^5x_4^{24}\\
& + x_2^3x_3x_4x_5^{30} + x_1^3x_3x_4x_5^{30} + x_1^3x_2x_4x_5^{30} + x_1^3x_2x_3x_5^{30} + x_1^3x_2x_3x_4^{30}\\
& + x_2^3x_3x_4^{30}x_5 + x_1^3x_3x_4^{30}x_5 + x_1^3x_2x_4^{30}x_5 + x_1^3x_2x_3^{30}x_5 + x_1^3x_2x_3^{30}x_4\\
& + x_2x_3x_4^6x_5^{27} + x_1x_3x_4^6x_5^{27} + x_1x_2x_4^6x_5^{27} + x_1x_2x_3^6x_5^{27} + x_1x_2x_3^6x_4^{27}\\
& + x_2x_3x_4^7x_5^{26} + x_1x_3x_4^7x_5^{26} + x_1x_2x_4^7x_5^{26} + x_1x_2x_3^7x_5^{26} + x_1x_2x_3^7x_4^{26}\\
& + x_2^3x_3^4x_4x_5^{27} + x_2^3x_3^4x_4^{27}x_5 + x_1^3x_3^4x_4x_5^{27} + x_1^3x_3^4x_4^{27}x_5 + x_1^3x_2^4x_4x_5^{27}\\
& + x_1^3x_2^4x_4^{27}x_5 + x_1^3x_2^4x_3x_5^{27} + x_1^3x_2^4x_3x_4^{27} + x_1^3x_2^4x_3^{27}x_5 + x_1^3x_2^4x_3^{27}x_4\\
& + x_2x_3^6x_4x_5^{27} + x_2x_3^6x_4^{27}x_5 + x_1x_3^6x_4x_5^{27} + x_1x_3^6x_4^{27}x_5 + x_1x_2^6x_4x_5^{27}
\end{aligned}$$

$$
\begin{aligned}
&+ x_1x_2^6x_4^{27}x_5 + x_1x_2^6x_3x_5^{27} + x_1x_2^6x_3x_4^{27} + x_1x_2^6x_3^{27}x_5 + x_1x_2^6x_3^{27}x_4\\
&+ x_2^3x_3x_4^4x_5^{27} + x_1^3x_3x_4^4x_5^{27} + x_1^3x_2x_4^4x_5^{27} + x_1^3x_2x_3^4x_5^{27} + x_1^3x_2x_3^4x_4^{27}\\
&+ x_2^3x_3x_4^7x_5^{24} + x_1^3x_3x_4^7x_5^{24} + x_1^3x_2x_4^7x_5^{24} + x_1^3x_2x_3^7x_5^{24} + x_1^3x_2x_3^7x_4^{24},
\end{aligned}
$$

$$
\begin{aligned}
p_5 = {} & x_2x_3^3x_4x_5^{30} + x_1x_3^3x_4x_5^{30} + x_1x_2^3x_4x_5^{30} + x_1x_2^3x_3x_5^{30} + x_1x_2^3x_3x_4^{30}\\
&+ x_2^3x_3^3x_4x_5^{28} + x_2^3x_3^3x_4^{28}x_5 + x_1^3x_3^3x_4x_5^{28} + x_1^3x_3^3x_4^{28}x_5 + x_1^3x_2^3x_4x_5^{28}\\
&+ x_1^3x_2^3x_4^{28}x_5 + x_1^3x_2^3x_3x_5^{28} + x_1^3x_2^3x_3x_4^{28} + x_1^3x_2^3x_3^{28}x_5 + x_1^3x_2^3x_3^{28}x_4\\
&+ x_2x_3^{30}x_4x_5^3 + x_2x_3^{30}x_4^3x_5 + x_1x_3^{30}x_4x_5^3 + x_1x_3^{30}x_4^3x_5 + x_1x_2^{30}x_4x_5^3\\
&+ x_1x_2^{30}x_4^3x_5 + x_1x_2^{30}x_3x_5^3 + x_1x_2^{30}x_3x_4^3 + x_1x_2^{30}x_3^3x_5 + x_1x_2^{30}x_3^3x_4\\
&+ x_2x_3^7x_4^{26}x_5 + x_2^7x_3x_4^{26}x_5 + x_1x_3^7x_4^{26}x_5 + x_1x_2^7x_4^{26}x_5 + x_1x_2^7x_3^{26}x_5\\
&+ x_1x_2^7x_3^{26}x_4 + x_1^7x_3x_4^{26}x_5 + x_1^7x_2x_4^{26}x_5 + x_1^7x_2x_3^{26}x_5 + x_1^7x_2x_3^{26}x_4\\
&+ x_2x_3^2x_4^3x_5^{29} + x_2x_3^2x_4^{29}x_5^3 + x_2x_3^3x_4^2x_5^{29} + x_2^3x_3x_4^2x_5^{29} + x_1x_3^2x_4^3x_5^{29}\\
&+ x_1x_3^2x_4^{29}x_5^3 + x_1x_3^3x_4^2x_5^{29} + x_1x_2^2x_4^3x_5^{29} + x_1x_2^2x_4^{29}x_5^3 + x_1x_2^2x_3^3x_5^{29}\\
&+ x_1x_2^2x_3^3x_4^{29} + x_1x_2^2x_3^{29}x_5^3 + x_1x_2^2x_3^{29}x_4^3 + x_1x_2^3x_4^2x_5^{29} + x_1x_2^3x_3^2x_5^{29}\\
&+ x_1x_2^3x_3^2x_4^{29} + x_1^3x_3x_4^2x_5^{29} + x_1^3x_2x_4^2x_5^{29} + x_1^3x_2x_3^2x_5^{29} + x_1^3x_2x_3^2x_4^{29}\\
&+ x_2x_3^2x_4^5x_5^{27} + x_1x_3^2x_4^5x_5^{27} + x_1x_2^2x_4^5x_5^{27} + x_1x_2^2x_3^5x_5^{27} + x_1x_2^2x_3^5x_4^{27}\\
&+ x_2x_3^2x_4^7x_5^{25} + x_2x_3^7x_4^2x_5^{25} + x_2^7x_3x_4^2x_5^{25} + x_1x_3^2x_4^7x_5^{25} + x_1x_3^7x_4^2x_5^{25}\\
&+ x_1x_2^2x_4^7x_5^{25} + x_1x_2^2x_3^7x_5^{25} + x_1x_2^2x_3^7x_4^{25} + x_1x_2^7x_4^2x_5^{25} + x_1x_2^7x_3^2x_5^{25}\\
&+ x_1x_2^7x_3^2x_4^{25} + x_1^7x_3x_4^2x_5^{25} + x_1^7x_2x_4^2x_5^{25} + x_1^7x_2x_3^2x_5^{25} + x_1^7x_2x_3^2x_4^{25}\\
&+ x_2x_3^3x_4^4x_5^{27} + x_1x_3^3x_4^4x_5^{27} + x_1x_2^3x_4^4x_5^{27} + x_1x_2^3x_3^4x_5^{27} + x_1x_2^3x_3^4x_4^{27}\\
&+ x_2x_3^3x_4^7x_5^{24} + x_2x_3^7x_4^3x_5^{24} + x_2^3x_3^7x_4x_5^{24} + x_2^7x_3x_4^3x_5^{24} + x_2^7x_3^3x_4x_5^{24}\\
&+ x_1x_3^3x_4^7x_5^{24} + x_1x_3^7x_4^3x_5^{24} + x_1x_2^3x_4^7x_5^{24} + x_1x_2^3x_3^7x_5^{24} + x_1x_2^3x_3^7x_4^{24}\\
&+ x_1x_2^7x_4^3x_5^{24} + x_1x_2^7x_3^3x_5^{24} + x_1x_2^7x_3^3x_4^{24} + x_1^3x_3^7x_4x_5^{24} + x_1^3x_2^7x_4x_5^{24}\\
&+ x_1^3x_2^7x_3x_5^{24} + x_1^3x_2^7x_3x_4^{24} + x_1^7x_3x_4^3x_5^{24} + x_1^7x_3^3x_4x_5^{24} + x_1^7x_2x_4^3x_5^{24}\\
&+ x_1^7x_2x_3^3x_5^{24} + x_1^7x_2x_3^3x_4^{24} + x_1^7x_2^3x_4x_5^{24} + x_1^7x_2^3x_3x_5^{24} + x_1^7x_2^3x_3x_4^{24}\\
&+ x_2x_3^3x_4^{28}x_5^3 + x_2^3x_3x_4^{28}x_5^3 + x_1x_3^3x_4^{28}x_5^3 + x_1x_2^3x_4^{28}x_5^3 + x_1x_2^3x_3^{28}x_5^3\\
&+ x_1x_2^3x_3^{28}x_4^3 + x_1^3x_3x_4^{28}x_5^3 + x_1^3x_2x_4^{28}x_5^3 + x_1^3x_2x_3^{28}x_5^3 + x_1^3x_2x_3^{28}x_4^3\\
&+ x_2^3x_3x_4^{30}x_5 + x_1^3x_3x_4^{30}x_5 + x_1^3x_2x_4^{30}x_5 + x_1^3x_2x_3^{30}x_5 + x_1^3x_2x_3^{30}x_4\\
&+ x_2x_3x_4^6x_5^{27} + x_1x_3x_4^6x_5^{27} + x_1x_2x_4^6x_5^{27} + x_1x_2x_3^6x_5^{27} + x_1x_2x_3^6x_4^{27}\\
&+ x_2x_3x_4^7x_5^{26} + x_1x_3x_4^7x_5^{26} + x_1x_2x_4^7x_5^{26} + x_1x_2x_3^7x_5^{26} + x_1x_2x_3^7x_4^{26}\\
&+ x_2^3x_3^4x_4x_5^{27} + x_2^3x_3^4x_4^{27}x_5 + x_1^3x_3^4x_4x_5^{27} + x_1^3x_3^4x_4^{27}x_5 + x_1^3x_2^4x_4x_5^{27}\\
&+ x_1^3x_2^4x_4^{27}x_5 + x_1^3x_2^4x_3x_5^{27} + x_1^3x_2^4x_3x_4^{27} + x_1^3x_2^4x_3^{27}x_5 + x_1^3x_2^4x_3^{27}x_4,
\end{aligned}
$$

$$\begin{aligned}p_6 = {} & x_2x_3^3x_4^{30}x_5 + x_1x_3^3x_4^{30}x_5 + x_1x_2^3x_4^{30}x_5 + x_1x_2^3x_3^{30}x_5 + x_1x_2^3x_3^{30}x_4\\
& + x_2x_3^{30}x_4x_5^3 + x_2x_3^{30}x_4^3x_5 + x_1x_3^{30}x_4x_5^3 + x_1x_3^{30}x_4^3x_5 + x_1x_2^{30}x_4x_5^3\\
& + x_1x_2^{30}x_4^3x_5 + x_1x_2^{30}x_3x_5^3 + x_1x_2^{30}x_3x_4^3 + x_1x_2^{30}x_3^3x_5 + x_1x_2^{30}x_3^3x_4\\
& + x_2x_3^7x_4^{26}x_5 + x_2^7x_3x_4^{26}x_5 + x_1x_3^7x_4^{26}x_5 + x_1x_2^7x_4^{26}x_5 + x_1x_2^7x_3^{26}x_5\\
& + x_1x_2^7x_3^{26}x_4 + x_1^7x_3x_4^{26}x_5 + x_1^7x_2x_4^{26}x_5 + x_1^7x_2x_3^{26}x_5 + x_1^7x_2x_3^{26}x_4\\
& + x_2x_3^2x_4^3x_5^{29} + x_2x_3^2x_4^{29}x_5^3 + x_2x_3^3x_4^2x_5^{29} + x_2^3x_3x_4^2x_5^{29} + x_1x_3^2x_4^3x_5^{29}\\
& + x_1x_3^2x_4^{29}x_5^3 + x_1x_3^3x_4^2x_5^{29} + x_1x_2^2x_4^3x_5^{29} + x_1x_2^2x_4^{29}x_5^3 + x_1x_2^2x_3^3x_5^{29}\\
& + x_1x_2^2x_3^3x_4^{29} + x_1x_2^2x_3^{29}x_5^3 + x_1x_2^2x_3^{29}x_4^3 + x_1x_2^3x_4^2x_5^{29} + x_1x_2^3x_3^2x_5^{29}\\
& + x_1x_2^3x_3^2x_4^{29} + x_1^3x_3x_4^2x_5^{29} + x_1^3x_2x_4^2x_5^{29} + x_1^3x_2x_3^2x_5^{29} + x_1^3x_2x_3^2x_4^{29}\\
& + x_2x_3^2x_4^5x_5^{27} + x_1x_3^2x_4^5x_5^{27} + x_1x_2^2x_4^5x_5^{27} + x_1x_2^2x_3^5x_5^{27} + x_1x_2^2x_3^5x_4^{27}\\
& + x_2x_3^2x_4^7x_5^{25} + x_2x_3^7x_4^2x_5^{25} + x_2^7x_3x_4^2x_5^{25} + x_1x_3^2x_4^7x_5^{25} + x_1x_3^7x_4^2x_5^{25}\\
& + x_1x_2^2x_4^7x_5^{25} + x_1x_2^2x_3^7x_5^{25} + x_1x_2^2x_3^7x_4^{25} + x_1x_2^7x_4^2x_5^{25} + x_1x_2^7x_3^2x_5^{25}\\
& + x_1x_2^7x_3^2x_4^{25} + x_1^7x_3x_4^2x_5^{25} + x_1^7x_2x_4^2x_5^{25} + x_1^7x_2x_3^2x_5^{25} + x_1^7x_2x_3^2x_4^{25}\\
& + x_2x_3^3x_4^4x_5^{27} + x_1x_3^3x_4^4x_5^{27} + x_1x_2^3x_4^4x_5^{27} + x_1x_2^3x_3^4x_5^{27} + x_1x_2^3x_3^4x_4^{27}\\
& + x_2x_3^3x_4^7x_5^{24} + x_2x_3^7x_4^3x_5^{24} + x_2^3x_3^7x_4x_5^{24} + x_2^7x_3x_4^3x_5^{24} + x_2^7x_3^3x_4x_5^{24}\\
& + x_1x_3^3x_4^7x_5^{24} + x_1x_3^7x_4^3x_5^{24} + x_1x_2^3x_4^7x_5^{24} + x_1x_2^3x_3^7x_5^{24} + x_1x_2^3x_3^7x_4^{24}\\
& + x_1x_2^7x_4^3x_5^{24} + x_1x_2^7x_3^3x_5^{24} + x_1x_2^7x_3^3x_4^{24} + x_1^3x_3^7x_4x_5^{24} + x_1^3x_2^7x_4x_5^{24}\\
& + x_1^3x_2^7x_3x_5^{24} + x_1^3x_2^7x_3x_4^{24} + x_1^7x_3x_4^3x_5^{24} + x_1^7x_3^3x_4x_5^{24} + x_1^7x_2x_4^3x_5^{24}\\
& + x_1^7x_2x_3^3x_5^{24} + x_1^7x_2x_3^3x_4^{24} + x_1^7x_2^3x_4x_5^{24} + x_1^7x_2^3x_3x_5^{24} + x_1^7x_2^3x_3x_4^{24}\\
& + x_2x_3^3x_4^{28}x_5^3 + x_2^3x_3x_4^{28}x_5^3 + x_1x_3^3x_4^{28}x_5^3 + x_1x_3^3x_4^{28}x_5^3 + x_1x_2^3x_3^{28}x_5^3\\
& + x_1x_2^3x_3^{28}x_4^3 + x_1^3x_3x_4^{28}x_5^3 + x_1^3x_2x_4^{28}x_5^3 + x_1^3x_2x_3^{28}x_5^3 + x_1^3x_2x_3^{28}x_4^3\\
& + x_2^3x_3x_4^{30}x_5 + x_1^3x_3x_4^{30}x_5 + x_1^3x_2x_4^{30}x_5 + x_1^3x_2x_3^{30}x_5 + x_1^3x_2x_3^{30}x_4\\
& + x_2x_3^6x_4x_5^{27} + x_2x_3^6x_4^{27}x_5 + x_1x_3^6x_4x_5^{27} + x_1x_3^6x_4^{27}x_5 + x_1x_2^6x_4x_5^{27}\\
& + x_1x_2^6x_4^{27}x_5 + x_1x_2^6x_3x_5^{27} + x_1x_2^6x_3x_4^{27} + x_1x_2^6x_3^{27}x_5 + x_1x_2^6x_3^{27}x_4\\
& + x_2^3x_3x_4^4x_5^{27} + x_1^3x_3x_4^4x_5^{27} + x_1^3x_2x_4^4x_5^{27} + x_1^3x_2x_3^4x_5^{27} + x_1^3x_2x_3^4x_4^{27}\\
& + x_2^3x_3x_4^7x_5^{24} + x_1^3x_3x_4^7x_5^{24} + x_1^3x_2x_4^7x_5^{24} + x_1^3x_2x_3^7x_5^{24} + x_1^3x_2x_3^7x_4^{24}.\end{aligned}$$

6.6.2. *The Σ_5-invariants of $QP_5^+(\omega_{(1)})$.*

$QP_5^+(\omega_{(1)})^{\Sigma_5} = \langle [p_7], [p_8], [p_9]\rangle$, where

$$\begin{aligned}p_7 = {} & x_1x_2x_3^6x_4x_5^{26} + x_1x_2x_3^6x_4^{26}x_5 + x_1x_2x_3^2x_4^5x_5^{26} + x_1x_2^2x_3x_4^5x_5^{26}\\
& + x_1x_2x_3^2x_4^6x_5^{25} + x_1x_2^2x_3x_4^6x_5^{25} + x_1x_2^2x_3^5x_4^2x_5^{25} + x_1x_2^2x_3^5x_4^{25}x_5^2\\
& + x_1x_2^3x_3^4x_4^2x_5^{25} + x_1x_2^3x_3^4x_4^{25}x_5^2 + x_1^3x_2x_3^4x_4^2x_5^{25} + x_1^3x_2x_3^4x_4^{25}x_5^2\\
& + x_1x_2^3x_3^5x_4^2x_5^{24} + x_1x_2^3x_3^5x_4^{24}x_5^2 + x_1^3x_2x_3^5x_4^2x_5^{24} + x_1^3x_2x_3^5x_4^{24}x_5^2\\
& + x_1x_2^3x_3^3x_4^4x_5^{24} + x_1^3x_2x_3^3x_4^4x_5^{24} + x_1^3x_2^3x_3x_4^4x_5^{24} + x_1^3x_2^3x_3^4x_4x_5^{24}\\
& + x_1^3x_2^3x_3^4x_4^{24}x_5 + x_1x_2^3x_2^2x_4x_5^{28} + x_1x_2^3x_3^2x_4^{28}x_5 + x_1^3x_2x_3^2x_4x_5^{28}\end{aligned}$$

$$
\begin{aligned}
&+ x_1^3x_2x_3^2x_4^{28}x_5 + x_1x_2^2x_3x_4^4x_5^{27} + x_1x_2^2x_3^4x_4x_5^{27} + x_1x_2^2x_3^4x_4^{27}x_5\\
&+ x_1x_2^2x_3x_4^7x_5^{24} + x_1x_2^2x_3^7x_4x_5^{24} + x_1x_2^2x_3^7x_4^{24}x_5 + x_1x_2^7x_3^2x_4x_5^{24}\\
&+ x_1x_2^7x_3^2x_4^{24}x_5 + x_1^7x_2x_3^2x_4x_5^{24} + x_1^7x_2x_3^2x_4^{24}x_5 + x_1x_2^2x_3^4x_4^9x_5^{19}\\
&+ x_1x_2^2x_3^4x_4^{11}x_5^{17} + x_1x_2^2x_3^5x_4^8x_5^{19} + x_1x_2^2x_3^5x_4^{11}x_5^{16} + x_1x_2^2x_3^7x_4^8x_5^{17}\\
&+ x_1x_2^7x_3^2x_4^8x_5^{17} + x_1^7x_2x_3^2x_4^8x_5^{17} + x_1x_2^2x_3^7x_4^9x_5^{16} + x_1x_2^7x_3^2x_4^9x_5^{16}\\
&+ x_1^7x_2x_3^2x_4^9x_5^{16} + x_1x_2^3x_3^4x_4^8x_5^{19} + x_1^3x_2x_3^4x_4^8x_5^{19} + x_1x_2^3x_3^4x_4^{11}x_5^{16}\\
&+ x_1^3x_2x_3^4x_4^{11}x_5^{16} + x_1x_2^3x_3^7x_4^8x_5^{16} + x_1x_2^7x_3^3x_4^8x_5^{16} + x_1^3x_2x_3^7x_4^8x_5^{16}\\
&+ x_1^3x_2^7x_3x_4^8x_5^{16} + x_1^7x_2x_3^3x_4^8x_5^{16} + x_1^7x_2^3x_3x_4^8x_5^{16},
\end{aligned}
$$

$$
\begin{aligned}
p_8 = {}& x_1x_2x_3^2x_4^3x_5^{28} + x_1x_2x_3^2x_4^{28}x_5^3 + x_1x_2x_3^3x_4^2x_5^{28} + x_1x_2^3x_3x_4^2x_5^{28}\\
&+ x_1^3x_2x_3x_4^2x_5^{28} + x_1x_2x_3^2x_4^4x_5^{27} + x_1x_2x_3^2x_4^7x_5^{24} + x_1x_2x_3^7x_4^2x_5^{24}\\
&+ x_1x_2^7x_3x_4^2x_5^{24} + x_1^7x_2x_3x_4^2x_5^{24} + x_1x_2^2x_3^3x_4^4x_5^{25} + x_1x_2^2x_3^4x_4^3x_5^{25}\\
&+ x_1x_2^2x_3^4x_4^{25}x_5^3 + x_1x_2^3x_3^2x_4^4x_5^{25} + x_1^3x_2x_3^2x_4^4x_5^{25} + x_1x_2^2x_3^3x_4^5x_5^{24}\\
&+ x_1x_2^2x_3^5x_4^3x_5^{24} + x_1x_2^2x_3^5x_4^{24}x_5^3 + x_1x_2^3x_3^2x_4^5x_5^{24} + x_1^3x_2x_3^2x_4^5x_5^{24}\\
&+ x_1x_2^3x_3^4x_4^3x_5^{24} + x_1x_2^3x_3^4x_4^{24}x_5^3 + x_1^3x_2x_3^4x_4^3x_5^{24} + x_1^3x_2x_3^4x_4^{24}x_5^3\\
&+ x_1x_2^2x_3^4x_4^9x_5^{19} + x_1x_2^2x_3^4x_4^{11}x_5^{17} + x_1x_2^2x_3^5x_4^8x_5^{19} + x_1x_2^2x_3^5x_4^{11}x_5^{16}\\
&+ x_1x_2^2x_3^7x_4^8x_5^{17} + x_1x_2^7x_3^2x_4^8x_5^{17} + x_1^7x_2x_3^2x_4^8x_5^{17} + x_1x_2^2x_3^7x_4^9x_5^{16}\\
&+ x_1x_2^7x_3^2x_4^9x_5^{16} + x_1^7x_2x_3^2x_4^9x_5^{16} + x_1x_2^3x_3^4x_4^8x_5^{19} + x_1^3x_2x_3^4x_4^8x_5^{19}\\
&+ x_1x_2^3x_3^4x_4^{11}x_5^{16} + x_1^3x_2x_3^4x_4^{11}x_5^{16} + x_1x_2^3x_3^7x_4^8x_5^{16} + x_1x_2^7x_3^3x_4^8x_5^{16}\\
&+ x_1^3x_2x_3^7x_4^8x_5^{16} + x_1^3x_2^7x_3x_4^8x_5^{16} + x_1^7x_2x_3^3x_4^8x_5^{16} + x_1^7x_2^3x_3x_4^8x_5^{16},
\end{aligned}
$$

$$
\begin{aligned}
p_9 = {}& x_1x_2^2x_3x_4^3x_5^{28} + x_1x_2^2x_3x_4^{28}x_5^3 + x_1x_2^2x_3^3x_4x_5^{28} + x_1x_2^2x_3^3x_4^{28}x_5\\
&+ x_1x_2^2x_3^{28}x_4x_5^3 + x_1x_2^2x_3^{28}x_4^3x_5 + x_1x_2^3x_3^2x_4x_5^{28} + x_1x_2^3x_3^2x_4^{28}x_5\\
&+ x_1^3x_2x_3^2x_4x_5^{28} + x_1^3x_2x_3^2x_4^{28}x_5 + x_1x_2^2x_3x_4^4x_5^{27} + x_1x_2^2x_3^4x_4x_5^{27}\\
&+ x_1x_2^2x_3^4x_4^{27}x_5 + x_1x_2^2x_3x_4^7x_5^{24} + x_1x_2^2x_3^7x_4x_5^{24} + x_1x_2^2x_3^7x_4^{24}x_5\\
&+ x_1x_2^7x_3^2x_4x_5^{24} + x_1x_2^7x_3^2x_4^{24}x_5 + x_1^7x_2x_3^2x_4x_5^{24} + x_1^7x_2x_3^2x_4^{24}x_5\\
&+ x_1x_2^2x_3^3x_4^4x_5^{25} + x_1x_2^2x_3^4x_4^3x_5^{25} + x_1x_2^2x_3^4x_4^{25}x_5^3 + x_1x_2^3x_3^2x_4^4x_5^{25}\\
&+ x_1^3x_2x_3^2x_4^4x_5^{25} + x_1x_2^2x_3^3x_4^5x_5^{24} + x_1x_2^2x_3^5x_4^3x_5^{24} + x_1x_2^2x_3^5x_4^{24}x_5^3\\
&+ x_1x_2^3x_3^2x_4^5x_5^{24} + x_1^3x_2x_3^2x_4^5x_5^{24} + x_1x_2^3x_3^4x_4^3x_5^{24} + x_1x_2^3x_3^4x_4^{24}x_5^3\\
&+ x_1^3x_2x_3^4x_4^3x_5^{24} + x_1^3x_2x_3^4x_4^{24}x_5^3 + x_1x_2^2x_3^4x_4^9x_5^{19} + x_1x_2^2x_3^4x_4^{11}x_5^{17}\\
&+ x_1x_2^2x_3^5x_4^8x_5^{19} + x_1x_2^2x_3^5x_4^{11}x_5^{16} + x_1x_2^2x_3^7x_4^8x_5^{17} + x_1x_2^7x_3^2x_4^8x_5^{17}\\
&+ x_1^7x_2x_3^2x_4^8x_5^{17} + x_1x_2^2x_3^7x_4^9x_5^{16} + x_1x_2^7x_3^2x_4^9x_5^{16} + x_1^7x_2x_3^2x_4^9x_5^{16}\\
&+ x_1x_2^3x_3^4x_4^8x_5^{19} + x_1^3x_2x_3^4x_4^8x_5^{19} + x_1x_2^3x_3^4x_4^{11}x_5^{16} + x_1^3x_2x_3^4x_4^{11}x_5^{16}\\
&+ x_1x_2^3x_3^7x_4^8x_5^{16} + x_1x_2^7x_3^3x_4^8x_5^{16} + x_1^3x_2x_3^7x_4^8x_5^{16} + x_1^3x_2^7x_3x_4^8x_5^{16}\\
&+ x_1^7x_2x_3^3x_4^8x_5^{16} + x_1^7x_2^3x_3x_4^8x_5^{16}.
\end{aligned}
$$

6.7. GL_5**-invariants of** $\mathbb{F}_2 \otimes_{\mathcal{A}} P_5$ **in degree** 15**.**

$(\mathbb{F}_2 \otimes_{\mathcal{A}} P_5)^{GL_5}_{15} = \langle [p], [q] \rangle$, where

$$\begin{aligned} p = {} & x_1^{15} + x_2^{15} + x_3^{15} + x_4^{15} + x_5^{15} + x_1x_2^{14} + x_1x_3^{14} + x_1x_4^{14} + x_1x_5^{14} \\ & + x_2x_3^{14} + x_2x_4^{14} + x_2x_5^{14} + x_3x_4^{14} + x_3x_5^{14} + x_4x_5^{14} + x_1x_2^2x_3^{12} \\ & + x_1x_2^2x_4^{12} + x_1x_2^2x_5^{12} + x_1x_3^2x_4^{12} + x_1x_3^2x_5^{12} + x_1x_4^2x_5^{12} + x_2x_3^2x_4^{12} \\ & + x_2x_3^2x_5^{12} + x_2x_4^2x_5^{12} + x_3x_4^2x_5^{12} + x_1x_2^2x_3^4x_4^8 + x_1x_2^2x_3^4x_5^8 \\ & + x_1x_2^2x_4^4x_5^8 + x_1x_3^2x_4^4x_5^8 + x_2x_3^2x_4^4x_5^8 + x_1x_2^2x_3^4x_4^4x_5^4, \end{aligned}$$

$$\begin{aligned} q = {} & x_1x_2x_3x_4^6x_5^6 + x_1x_2x_3^6x_4x_5^6 + x_1x_2x_3^6x_4^6x_5 + x_1x_2^6x_3x_4x_5^6 \\ & + x_1x_2^6x_3x_4^6x_5 + x_1x_2^3x_3^6x_4x_5^4 + x_1x_2^3x_3^6x_4^4x_5 + x_1x_2^6x_3^3x_4x_5^4 \\ & + x_1x_2^6x_3^3x_4^4x_5 + x_1^3x_2x_3x_4^4x_5^6 + x_1^3x_2x_3x_4^6x_5^4 + x_1^3x_2x_3^4x_4x_5^6 \\ & + x_1^3x_2x_3^4x_4^6x_5 + x_1^3x_2^4x_3x_4x_5^6 + x_1^3x_2^4x_3x_4^6x_5 + x_1x_2^3x_3^3x_4^4x_5^4 \\ & + x_1^3x_2x_3^3x_4^4x_5^4 + x_1^3x_2^3x_3x_4^4x_5^4 + x_1^3x_2^3x_3^4x_4x_5^4 + x_1^3x_2^3x_3^4x_4^4x_5 \\ & + x_1^3x_2^4x_3^3x_4x_5^4 + x_1^3x_2^4x_3^3x_4^4x_5. \end{aligned}$$

Acknowledgment. The author would like to thank Prof. Nguyễn H. V. Hưng for valuable discussions on Kameko's squaring operation.

The first version of this paper was written when the author was visiting Viet Nam Institute for Advanced Study in Mathematics (VIASM). He would like to thank the VIASM for supporting the visit and kind hospitality.

References

[1] J. M. Boardman, *Modular representations on the homology of power of real projective space*, in: M. C. Tangora (Ed.), Algebraic Topology, Oaxtepec, 1991, in: Contemp. Math., vol. 146, 1993, pp. 49-70. MR1224907

[2] R. R. Bruner, L. M. Hà, N. H. V. Hưng, *On behavior of the algebraic transfer*, Trans. Amer. Math. Soc. 357 (2005), 473-487. MR2095619

[3] D. P. Carlisle, R. M. W. Wood, *The boundedness conjecture for the action of the Steenrod algebra on polynomials*, in: N. Ray and G. Walker (ed.), Adams Memorial Symposium on Algebraic Topology 2, (Manchester, 1990), in: London Math. Soc. Lecture Notes Ser., Cambridge Univ. Press, Cambridge, vol. 176, 1992, pp. 203-216. MR1232207

[4] T. W. Chen, *Determination of* $\mathrm{Ext}^{5,*}_{\mathcal{A}}(\mathbb{Z}/2, \mathbb{Z}/2)$, Topology Appl., 158 (2011), 660-689. MR2774051

[5] P. H. Chơn, L. M. Hà, *Lambda algebra and the Singer transfer*, C. R. Acad. Sci. Paris Ser. I 349 (2011), 21-23. MR2755689

[6] P. H. Chơn, L. M. Hà, *On May spectral sequence and the algebraic transfer*, Manuscripta Math. 138 (2012), 141-160. MR2898751

[7] P. H. Chơn, L. M. Hà, *On the May spectral sequence and the algebraic transfer II*, Topology Appl. 178 (2014), 372-383. MR3276753

[8] L. M. Hà, *Sub-Hopf algebras of the Steenrod algebra and the Singer transfer*, "Proceedings of the International School and Conference in Algebraic Topology, Hà Nội 2004", Geom. Topol. Monogr., Geom. Topol. Publ., Coventry, vol. 11 (2007), 81-105. MR2402802

[9] N. H. V. Hưng, *The weak conjecture on spherical classes*, Math. Zeit. 231 (1999), 727-743. MR1709493

[10] N. H. V. Hưng, *The cohomology of the Steenrod algebra and representations of the general linear groups*, Trans. Amer. Math. Soc. 357 (2005), 4065-4089. MR2159700
[11] N. H. V. Hưng, V. T. N. Quỳnh, *The image of Singer's fourth transfer,* C. R. Acad. Sci. Paris Ser. I 347 (2009), 1415-1418. MR2588792
[12] M. Kameko, *Products of projective spaces as Steenrod modules*, PhD Thesis, The Johns Hopkins University, ProQuest LLC, Ann Arbor, MI, 1990, 29 pp. MR2638633
[13] W. H. Lin, $\mathrm{Ext}_{\mathcal{A}}^{4,*}(\mathbb{Z}/2,\mathbb{Z}/2)$ *and* $\mathrm{Ext}_{\mathcal{A}}^{5,*}(\mathbb{Z}/2,\mathbb{Z}/2)$, Topology Appl., 155 (2008), 459-496. MR2380930
[14] N. Minami, *The iterated transfer analogue of the new doomsday conjecture*, Trans. Amer. Math. Soc. 351 (1999), 2325-2351. MR1443884
[15] T. N. Nam, *Transfert algébrique et action du groupe linéaire sur les puissances divisées modulo 2*, Ann. Inst. Fourier (Grenoble) 58 (2008), 1785-1837. MR2445834
[16] F. P. Peterson, *Generators of* $H^*(\mathbb{R}P^\infty \times \mathbb{R}P^\infty)$ *as a module over the Steenrod algebra*, Abstracts Amer. Math. Soc. No. 833 (1987), 55-89.
[17] Đ. V. Phúc, N. Sum, *On the generators of the polynomial algebra as a module over the Steenrod algebra*, C. R. Acad. Sci. Paris, Ser. I 353 (2015), 1035-1040. MR3419856
[18] V. T. N. Quỳnh, *On behavior of the fifth algebraic transfer*, "Proceedings of the International School and Conference in Algebraic Topology, Hà Nội 2004", Geom. Topol. Monogr., Geom. Topol. Publ., Coventry, vol. 11 (2007), 309-326. MR2402811
[19] W. M. Singer, *The transfer in homological algebra*, Math. Zeit. 202 (1989), 493-523. MR1022818
[20] W. M. Singer, *On the action of the Steenrod squares on polynomial algebras*, Proc. Amer. Math. Soc. 111 (1991), 577-583. MR1045150
[21] N. E. Steenrod, D. B. A. Epstein, *Cohomology operations*, Annals of Mathematics Studies 50, Princeton University Press, Princeton N.J (1962). MR0145525
[22] N. Sum, *The negative answer to Kameko's conjecture on the hit problem*, Adv. Math. 225 (2010), 2365-2390. MR2680169
[23] N. Sum, *On the Peterson hit problem of five variables and its applications to the fifth Singer transfer*, East-West J. of Mathematics, 16 (2014), No. 1, 47-62. MR3409252
[24] N. Sum, *On the Peterson hit problem*, Adv. Math. 274 (2015), 432-489. MR3318156
[25] M. C. Tangora, *On the cohomology of the Steenrod algebra*, Math. Zeit. 116 (1970), 18-64. MR0266205
[26] N. K. Tín, N. Sum, *Kameko's homomorphism and the algebraic transfer*, C. R. Acad. Sci. Paris Ser. I 354 (2016), 940-943, MR3535350.
[27] N. K. Tín, *On Singer's conjecture for the fifth algebraic transfer*, Preprint 2016, available online at https://arxiv.org/abs/1609.02250.
[28] R. M. W. Wood, *Steenrod squares of polynomials and the Peterson conjecture*, Math. Proc. Cambridge Phil. Soc. 105 (1989), 307-309. MR0974986

Department of Mathematics, Quy Nhơn University,
170 An Dương Vương, Quy Nhơn, Bình Định, Viet Nam.
E-mail: nguyensum@qnu.edu.vn